This is a pre- or post-print of an article published in
Wardecki, T., Brötz, E., De Ford, C., Von Loewenich,
F.D., Rebets, Y., Tokovenko, B., Luzhetskyy, A., Merfort,
I.
Endophytic *Streptomyces* in the traditional medicinal
plant *Arnica montana* L.: Secondary metabolites and
biological activity
(2015) Antonie van Leeuwenhoek, International Journal of
General and Molecular Microbiology, 108 (2), art. no.
A012, pp. 391-402.

**Endophytic *Streptomyces* in the traditional medicinal plant
Arnica montana L.: Secondary metabolites and biological
activity**

Journal: Antonie van Leeuwenhoek

Tina Wardecki¹, Elke Brötz², Christian De Ford^{1,3}, Friederike D. von
Loewenich^{4*}, Yuriy Rebets², Bogdan Tokovenko², Andriy Luzhetskyy², Irmgard
Merfort¹

¹ Pharmaceutical Biology and Biotechnology, Albert-Ludwigs-University
Freiburg, 79104 Freiburg, Germany

² Helmholtz Institute for Pharmaceutical Research Saarland, Universität des
Saarlandes, 66123 Saarbrücken, Germany

³ Spemann Graduate School of Biology and Medicine (SGBM), Albert-Ludwigs-
University Freiburg, 79104 Freiburg, Germany

⁴ Institute for Medical Microbiology and Hygiene, Medical Center - University of
Freiburg, 79104 Freiburg, Germany

*present address: Department of Medical Microbiology and Hygiene, University
of Mainz -Medical Centre, 55131 Mainz, Germany

Corresponding author:

Prof. Dr. Irmgard Merfort

Pharmaceutical Biology and Biotechnology

Stefan-Meier-Strasse 19

79104 Freiburg, Germany

e-mail: Irmgard.Merfort@pharmazie.uni-freiburg.de

fax: + 49 (0) 761 / 203 8383

phone: + 49 (0) 761 / 203 8373

ABSTRACT

Arnica montana L. is a medical plant of the Asteraceae family and grows preferably on nutrient poor soils in mountainous environment. Such surroundings are known to make plants dependent on symbiosis with other organisms. Up to now only arbuscular mycorrhizal fungi were found to act as endophytic symbiosis partners for *A. montana*. Here we identified five *Streptomyces* strains, microorganisms also known to occur endophytic in plants and to produce a huge variety of active secondary metabolites, as inhabitants of *A. montana*. The secondary metabolite spectrum of these strains does not contain sesquiterpene lactones, but consists of the glutarimide antibiotics cycloheximide and actiphenol as well as the diketopiperazines cyclo-prolyl-valyl, cyclo-prolyl-isoleucyl, cyclo-prolyl-leucyl and cyclo-prolyl-phenylalanyl. Exemplarily, genome analysis of one strain was performed and indicated a huge genome size with a high number of natural products gene clusters among which genes for cycloheximide production were detected. Only weak activity against the gram-positive bacterium *Staphylococcus aureus* was revealed, but the extracts showed a marked cytotoxic activity as well as an antifungal activity against *Candida parapsilosis* and *Fusarium verticillioides*. Altogether, our results provide evidence that *A. montana* and its endophytic *Streptomyces* benefit from each other by completing their protection against competitors and pathogens and by exchanging plant growth promoting signals against nutrients.

ABBREVIATIONS

Act D, Actinomycin D, ANOVA, analysis of variance; EUCAST, European Committee on Antimicrobial Susceptibility Testing; ESCOP, European Scientific

Cooperative on Phytotherapy; HRESI, High-resolution electrospray ionization; NIST, National Institute of Standards and Technology; UPLC, Ultra Performance Liquid Chromatography.

KEYWORDS

natural products, symbiotic plant bacteria, *Arnica* root, cytotoxic and antifungal activity

INTRODUCTION

Arnica montana L. is a herbaceous plant of the Asteraceae family. It is endemic in Europe and predominantly found in nutrient-poor grasslands, shrublands, and in alpine mountain environments (Sugier et al. 2013). Preparations, especially from its flowerheads have a long lasting tradition for the external treatment of inflammatory diseases. They have got a positive judgment from the European Scientific Cooperative on Phytotherapy (ESCOP) and the European Pharmacopoeia for the treatment of several injuries such as bruises, distortions, sprains, hematoma, inflamed insect stings as well as rheumatic complaints (ESCOP 2003; Ph.Eur. 2012). Sesquiterpene lactones such as helenalin, 11 α ,13-dihydrohelenalin and their esters are the most effective compounds responsible for the utilised anti-inflammatory activity (Teuscher et al. 2009; Merfort 2010). These compounds were proven to inhibit the transcription factor NF- κ B thus targeting the inflammatory processes at a central point (Lyss et al. 1997; García-Piñeres et al. 2001). Additionally, those molecules exert an antibacterial activity against gram-positive bacteria (Lee et al. 1977).

Even though *A. montana* grows on nutrient-poor acidic soils, which makes

plants often more dependent on symbiosis with other organisms, only few details about symbiotic interactions with *A. montana* are known. There is evidence for symbiosis with arbuscular mycorrhizal fungi, but no report exists on a symbiosis with endophytic *Streptomyces* (Ryszka et al. 2010). Up to now only a few species of the large plant family of Asteraceae were reported to interact with endophytic *Streptomyces*, among them *Artemisia annua*, which is also a plant with known medicinal properties (Sardi et al. 1992; Jiménez-Esquilín and Roane 2005; Castillo et al. 2007; Lin et al. 2012; Tanvir et al. 2014).

Streptomyces and other endophytes are known to produce various bioactive metabolites which target e.g. bacteria, fungi, and cancer cells (Porrás-Alfaro and Bayman 2011; Bérdy 2005). Regarding the secondary metabolites of plants, it is known, that in several cases the real producer of the compounds is not the plant itself but its endophytes (Bérdy 2005). There are also cases in which both, plant and endophyte produce the same metabolites. One prominent example is the anticancer drug taxol which is isolated from the bark of *Taxus brevifolia*. It has been shown that the inhabiting endophyte *Taxomyces andreanae* is also able to synthesize this substance (Stierle et al. 1993). This arises the question if endophytes of *A. montana* are also able to produce the medically active sesquiterpene lactone derivatives synthesised by this plant.

The aims of this study were to identify species of *Streptomyces* inhabiting *A. montana*, as well as to determine active secondary metabolites in the *Streptomyces* extracts. From the possible huge variety of pharmacological effects, we focused on the cytotoxic, antibacterial and antifungal properties of these extracts. Altogether these studies will further increase the knowledge about this medicinal plant and its interactions with microorganisms.

METHODS

Plant material and isolation of bacterial strains

A. montana L. was collected in Vosges mountains, Breitfirstkopf, in France 2009. The isolation of actinobacteria was performed as described in literature (Coombs and Franco 2003) with minor modifications. *A. montana* L. roots from three plants were dried for 48 h at room temperature, and were afterwards thoroughly washed to remove all soil rests. The roots were excised into pieces with a diameter of approximately 0.3-0.7 cm and a length of 2-3 cm. This material was subjected to a three-step surface sterilisation procedure: a 60seconds wash in 99% ethanol, followed by a 6-minute wash in 3.125% NaOCl, a 30-seconds wash in 99% ethanol, and a final rinse with sterile water. The surface-sterilised roots were then smashed into small fragments under sterile conditions and plated onto the isolation media at 28°C for up to 2 weeks. TSB and MS agar media were used in the isolation experiment. Finally, the strains were isolated on the MS agar, because on TSB agar medium *Actinomycetes* were overgrown by other bacteria. The efficacy of the surface sterilisation procedure was assessed by rolling surface-sterilised roots in the isolation medium which was then incubated at 28°C. Endophytic bacterial strains, in total 5, were stored in 20 % saccharose solution at -80 °C.

Cultivation of bacterial strains and preparation of the extracts

The bacterial strains were pre-cultivated in 300 mL Erlenmeyer flasks (2 days, 28 °C, 180 rpm) filled with 50 mL TSB medium and 100 µL of sucrose-culture (1 L TSB consists of tryptic soy broth 30.0 g). The main cultivation was made in 500 mL Erlenmeyer flasks containing 100 mL DNPM-medium (1 litre DNPM medium consists of dextrin 40.0 g, soy tone 7.50 g, fresh yeast 5.00 g, 4-

morpholinopropanesulfonic acid (MOPS) 21.0 g, pH 6.8). The cultures were inoculated with 5 mL pre-culture and cultivated for 7 days (30 °C, 180 rpm). Altogether five extracts were obtained. For confirmation of the presence of diketopiperazines NL-5 medium was used instead of DNPM medium (1 litre NL-5 medium consists of NaCl 1.00 g, K₂HPO₄ 1.00 g, MgSO₄ × 7 H₂O 0.50 g, glycerine 25.0 g, l-glutamine 5.84 g, trace-element-solution 2.00 mL, pH 7.3; trace-element-solution consists of ZnCl 40.0 mg, FeCl₃ × 6 H₂O 200 mg, CuCl₂ × 2 H₂O 10.0 mg, MnCl₂ × 4 H₂O, Na₂B₄O₇ × 10 H₂O 10.0 mg, (NH₄)Mo₇O₂₄ × 4 H₂O). For extraction the whole 100 mL cultures were stirred with 100 mL ethyl acetate for 30 min and were subsequently centrifuged (4000 rpm, 9 min) to separate biomass and aqueous phase from organic layer. The organic layer was concentrated under reduced pressure, transferred into 10 mL glass vials and evaporated using a sample concentrator with a light nitrogen flow and 40°C. The samples were re-suspended in 500 µL (200 fold concentration) methanol.

Analysis of bacterial metabolites

Methanolic extracts were analysed by GC-MS (GC-system: Agilent 6890 Series; detector: Agilent 5973 Network Mass Selective Detector, Agilent Technologies; ionization: electron ionization with 70 eV ionization energy) using a fused silica capillary column (30 m x 0.25 mm) coated with 0.25 µm dimethyl polysiloxane (RTX®-1 MS, Restek) and helium as carrier gas with a flow rate of 1.0 mL per minute. The temperature profile started at 120 °C followed by a heating rate of 10 °C per minute to 270 °C, which was maintained for 15 minutes. The injector and detector temperatures were 280 °C, the injection volume was 1.0 µL and was split 10:1. Obtained mass spectra were compared with National Institute of

Standards and Technology (NIST) 11 reference database and a self-assembled database of sesquiterpene lactones and their esters found in *A. montana*. This comparison led to propositions for molecules with the best matching fragmentation pattern. Further confirmation was done by comparing the mass spectra with those available in literature (for actiphenol and diketopiperazines) or obtained with purchased reference substance (for cycloheximide, Sigma-Aldrich). Dereplication of results was conducted by UPLC-UV-HRESI-MS-system at LTQ Orbitrap (Thermo Scientific) using 18 min gradient starting with solvent B from 5% up to 95% (A: water + 0.1% formic acid, B: CH₃CN + 0.1% formic acid) on waters BEH C18 column (100 × 2.1, 1.7 µm). The flow was set to 0.6 mL per minute, column temperature to 45°C and injection to 1 µL per sample). MS and UV data were analysed using DNP (Dictionary of Natural Products, CRD Press, Frances and Taylor).

Cell culture

Jurkat T cells obtained from DSMZ (Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH) were cultivated in Roswell Park Memorial Institute (RPMI) 1640 medium supplemented by 10 % heat-inactivated foetal bovine serum (both Gibco®) and 1 mL penicillin/streptomycin solution (Roche Diagnostics). Cells were grown in a humidified atmosphere at 37 °C with 5 % CO₂. Medium was renewed regularly.

MTT assay

For this colorimetric cell viability assay from Mosmann (1983) Jurkat T cells were seeded in 96 well plates (15.000 cells per well in 150 µL RPMI 1640

medium). Subsequently, the cells were incubated overnight in a humidified atmosphere at 37 °C with 5 % CO₂. On the following day, the cells were incubated for 24 h with 10 or 50 µg of the *Streptomyces* extracts mL⁻¹ dissolved in DMSO. For each condition three technical replicates were conducted. In the last two hours of the incubation time the cells were additionally treated with 0.5 mg 3-[4,5-dimethylthiazol-2-yl]-2,5-diphenyltetrazolium bromide (MTT, Roth) mL⁻¹ in phosphate buffered saline (PBS). The obtained formazan crystals were dissolved in DMSO and the absorption of the solution was measured by a microplate reader (Bio-Rad) at 595 nm.

The values of the samples were referred to untreated cells as control. Viability was calculated as percentage of surviving cells compared with the mentioned untreated cells. 0.4 µg Actinomycin D mL⁻¹ (Axxora) was used as positive control and 0.25 % DMSO as negative control. This viability test was performed twice.

Analysis of antibacterial activity

Antimicrobial susceptibility testing was done by disc diffusion test using the reproducible method of EUCAST (Matuschek et al. 2014). Three gram-positive strains (*Staphylococcus aureus* (ATCC 29213), *Enterococcus faecalis* (ATCC 29212), *Streptococcus pneumoniae* (ATCC 49619) and three gram-negative bacterial strains (*Escherichia coli* (ATCC 25922), *Pseudomonas aeruginosa* (ATCC 27583), *Haemophilus influenzae* (NCTC 8468) were tested. According to EUCAST guidelines the inoculum suspensions were adjusted to the density of a 0.5 McFarland standard. The inoculum suspension was then regularly spread over the agar plate using an automatic plate rotator. For *S. aureus*, *E. faecalis*, *E. coli*, and *P. aeruginosa* and Mueller-Hinton agar without

supplements was utilised. Those strains were incubated at 35 °C for 16 – 20 hours after adding the paper disks. Aberrantly *S. pneumoniae* and *H. influenzae* were grown on Mueller-Hinton agar with 5 % mechanically defibrinated horse blood and 20 mg β -nicotinamide adenine dinucleotide L⁻¹ and kept at 35 °C with 5 % CO₂ for 16 – 20 hours after adding the testing disks.

The *Streptomyces* extracts were tested in concentrations of 10, 50, and 100 μ g per disk. As negative control the solvent of the extracts, methanol, was applied in the highest used volume of 20 μ L per disc. Different positive controls were used for the different strains, *E. coli* susceptibility with 10 μ g of ampicillin, *S. aureus* and *S. pneumoniae* with 1 unit of penicillin, *P. aeruginosa* with 30 μ g piperacillin, *E. faecalis* and *H. influenzae* with 2 μ g ampicillin (all from BD Sensi-Disc™). In the case of extract 10 tested against *S. aureus* nalidixic acid (Sigma-Aldrich) was also analysed using concentrations of 15 and 30 μ g. After incubation the zone diameter of the inhibition zones was measured with a calliper. This disc diffusion test was performed at least two times.

Analysis of antifungal properties

Extracts 2, 3, and 4 were tested for their effects on the fungi *Candida parapsilosis* (DSM 5784) and *Fusarium verticillioides* (DSM 62264) using the disc diffusion method. These fungal strains were grown in yeast medium (1 litre of medium consists of 3 g yeast extract, 3 g malt extract, 5 g peptone from soybean and 10 g glucose). To prepare agar plates 20 g agar was used for one litre of yeast medium. Methanol was used as negative control and 75 μ g of nystatin as positive control. The inoculated plates were kept at 28 °C for 16 – 20 (*C. parapsilosis*) resp. 24 – 48 (*F. verticillioides*) hours before evaluation. The disc diffusion test was performed two times.

Genome sequencing and analysis

Genome sequencing of the cultivated strain was performed using Illumina MiSEQ technology. High molecular mass DNA was extracted from the selected strain, and two libraries were created: 8kb mate-pair library, as well as a shorter-insert (~600bp) shotgun paired-end library. Reads of an approximate length 2x250 (before trimming) were obtained for both libraries. Newbler software version 2.8 was used for genome assembly.

In the first step, gene finding was done using GISMO (Krause et al. 2007) followed by GenDB 2.0 automatic annotation (Meyer et al. 2003). In the second annotation step, all predicted ORFs were manually re-inspected to correct start codon and function assignments. For the identification of secondary metabolite clusters, the genome of *Streptomyces* sp. Arnica3 was scanned for homologues to known secondary metabolite synthases via BLAST search. These manual investigations were supported by antiSMASH (Medema et al. 2011).

16sDNA sequencing and phylogenetic analysis

Isolated strains 3 and 5 were grown in 10 ml of TSB media at 28°C for 3 days and total DNA was isolated as described (Kieser et al. 2000). Amplification of the 16S rRNA gene was carried out with primers: 8F (AGAGTTTGATYMTGGCTCAG) and 1510R (TACGGYTACCTTGTTACGACTT). Obtained fragments were gel purified using QIAquick Gel Extraction Kit (Qiagen, Venlo, Netherlands) and sequenced with the amplification primers 8F and 1510R generating almost the entire gene sequence (1365~1401 bp). 31 rDNA sequences were aligned using MAFFT

v7.017 (gap open penalty 1.53, offset value 0.123, scoring matrix 200PAM/k=2, algorithm: auto) (Kato et al. 2002). The dendrogram was built using Geneious (Kearse et al. 2012). The following settings were chosen: Tamura-Nei genetic distance model, neighbor-joining method, *E. coli* as an outgroup, bootstrap value 1000, consensus tree with 50 % support threshold. Tree branches were then proportionally transformed for display. Ribosomal RNA sequences for strains were extracted from genome annotations in GenBank. Based on the 16S phylogeny of the two isolated *Streptomyces* sp. (Fig. 4), they are the most similar ones to *S. avidinii* and *S. spororaveus*, followed by *S. fulvissimus* and *S. griseus subsp. griseus*.

Statistical analysis

Statistical analysis was performed with GraphPad Prism 5 software (GraphPad Software Inc.). Values are shown as means $\pm$ standard deviation (SD). For comparison between groups analysis of variance (ANOVA) followed by Bonferroni's post-test was used. *p*-values <0.05 were considered as significant.

RESULTS

Identification of secondary metabolites

The five extracts from the isolated actinobacteria were analysed by GC-MS and LC-MS. The diketopiperazines cyclo-prolyl-valyl, cyclo-leucyl-prolyl, and cyclo-isoleucyl-prolyl could be identified in all five extracts, whereas cycloheximide, actiphenol, and cyclo-prolyl-phenylalanyl only occurred in some of the extracts (see Table 1, for structures see Fig. 1). Cyclo-isoleucyl-prolyl and cyclo-prolyl-

valyl were not present in the NIST11 database. Identification was done by comparison with published fragmentation patterns (Bitzer 2005; Wang et al. 2010; Ortiz-Castro et al. 2011). Only extract 3 contained all mentioned molecules and the respective strain was therefore chosen for genome sequencing.

All found compounds were reported in literature as metabolites produced by *Streptomyces* (Highet and Prelog 1959; Macherla et al. 2005; Pettit et al. 2006). Nevertheless, diketopiperazines are also known to exist in autoclaved peptone containing medium used for cultivation of *Streptomyces* (Tamura et al. 1964). This can be explained by non-enzymatic cyclisation of linear dipeptides at extreme temperatures (Skwierczynski and Connors 1993). Due to the usage of DNPM medium with soytone as amino acid source it was necessary to conduct further experiments to confirm the diketopiperazines as metabolites from *Streptomyces*. On the one hand a mock cultivation of DNPM medium without adding *Streptomyces* was performed, on the other hand the *Streptomyces* strains were cultivated in NL-5 medium, whose only amino acid is glutamine. The yielded extracts were analysed and revealed the presence of diketopiperazines after cultivation in NL-5 as well as in the mock extracts of DNPM medium. Therefore, it can be concluded that the identified diketopiperazines are present as artefact from DNPM medium but also as secondary metabolites produced by *Streptomyces*.

Viability reducing properties of the *Streptomyces* extracts

Due to their known cytotoxic properties (Otani et al. 1989; Lakhvich et al. 1993), the identification of cycloheximide and actiphenol in some of the *Streptomyces* extracts lead to the assumption that those extracts could have an impact on the

viability of cancer cells. Therefore, we performed MTT assays using Jurkat T cells, an immortalised human T lymphocyte cell line. The cells were treated with either 0.4 $\mu\text{g mL}^{-1}$ actinomycin D as positive control, 0.25 % DMSO as negative control or with the *Streptomyces* extracts at 10 or 50 $\mu\text{g mL}^{-1}$ concentrations. Extracts 2, 3, 4, and 5 exhibited a significant and concentration dependent reduction of cell viability (see Fig. 2). In contrast, no reduction of viability was observed with extract 1. In summary, all extracts containing either cycloheximide or actiphenol or even both molecules showed significant viability reducing effects against Jurkat T cells.

Furthermore a caspase-3/-7 assay was performed for the most cytotoxic extracts 2 and 3 to determine whether the occurring cell death is caused by apoptosis (data not shown). The test showed no marked increase in caspase-3/-7 activity, indicating that this mechanism of cell death does not play any role.

Antibacterial activity of the *Streptomyces* extracts

Antibacterial properties are often found for metabolites from *Streptomyces*. Therefore, we investigated the effect of the extracts on six common bacterial test strains. Because three gram-positive and also three gram-negative strains with varying characteristics were used, this would provide a reasonable overview on the antibacterial activity of the examined extracts. Only extract 4 showed a slight concentration dependent activity against *S. aureus*, one of the gram-positive bacterial strains. This effect manifested itself in a zone around the test disks with a diffuse decline in bacterial population compared to the rest of the inoculated and incubated agar plate. The same situation could be shown using 15 μg of nalidixic acid, to which *S. aureus* is resistant (Ball 2000). In contrast, 1 unit of penicillin leads to a more definite inhibition zone (Fig. 3).

Hence it can be stated that there is a slight activity of extract 4 against *S. aureus*, but that the presence of colonies of *S. aureus* in the inhibition zone may imply a development of resistance as it is the case for nalidixic acid.

Additionally to the *Streptomyces* extracts, 100 µg of the mock extract containing DNPM medium without *Streptomyces*, which also comprised diketopiperazines, was tested against *S. aureus*. No zone of inhibition could be detected.

Antifungal activity of the *Streptomyces* extracts

Cycloheximide is known for its antifungal effect (Sharma et al. 2010). Therefore, the cycloheximide containing extracts 2 and 3 were tested for their antifungal properties in comparison with cycloheximide-lacking extract 4. As test strains the pathogenic yeast *Candida parapsilosis*, which can be found in soil, and the mold fungus *Fusarium verticillioides*, a fungus known to inhabit soil and plants, were used (Brown et al. 2006; Nosek et al. 2009). As presumed extracts 2 and 3 but not 4 showed growth-inhibitory activity against *Candida parapsilosis* and *Fusarium verticillioides* (Table 2).

Genome analysis

One of the isolated *Streptomyces* strains producing extract 3 was sequenced as described in Materials and Methods, resulting in 7 contigs, the longest of which (9206603 bp) contains 96% of all the sequence data and is one of the largest sequenced *Streptomyces* genomes. Judging by the genome size, this *Streptomyces* sp. is probably well-fit for highly-variable nutrient and environmental conditions. This genome contains 8319 predicted protein coding sequences (yielding genome coding density of 85.7%), 87 tRNAs, 1 tmRNA, 23 other RNAs, and 39 putative secondary metabolite clusters. GC content of this

genome is 70.6%, which is common for *Streptomyces*. Of the 39 putative secondary metabolite clusters, 1 was identified as siderophore, 1 as ectoine, 7 as terpene, 8 (2 per type) of bacteriocin, butyrolactone, lantipeptide and melanin, 9 NRPS, 4 PKS, and 9 clusters of other and hybrid types.

Finally, we decided to check if the sequenced strain has genes for cycloheximide production. We have used cycloheximide biosynthesis cluster sequence from GenBank, accession number JX014302.1 (*Streptomyces* sp. YIM 56141). We succeeded in finding all 10 ChxA-ChxJ genes. Average gene-level nucleotide identity to JX014302.1 is 79% with sequence coverage in 92-100% range (average 97%). In addition, we have performed the phylogenetic analysis of two isolated strains using 16S ribosomal RNA encoding genes. Based on the 16S phylogeny of the two isolated *Streptomyces* sp. are evolutionary closed to each other (Fig. 4), and they are the most similar to *S. avidinii* and *S. spororaveus*, followed by *S. fulvissimus* and *S. griseus* subsp. *griseus*.

DISCUSSION

Streptomyces produce a huge variety of bioactive molecules, including approximately 55 % of all clinically useful antibiotic drugs (Demain 1999). Biosynthesis of secondary metabolites in *Streptomyces* is dependent on manifold environmental criteria. Those criteria include temperature, light, pH, concentration of phosphate and oxygen and the amount and kind of carbon and nitrogen source (Ruiz et al. 2010). Therefore it is attractive to examine those bacterial species inhabiting environments with unusual ecological niches. One

of those ecological niches is occupied by *A. montana*, a plant which survives in a nutrient-poor hostile environment. Soil is known to be a harsh, competitive environment for microorganisms revealing extreme temperatures, moisture, and poor nutrient (Loria et al. 2006). This pertains especially in the mountainous environment without the massive usage of fertilizers, where *A. montana* is growing preferably (Sugier et al. 2013). It is not surprising that *Streptomyces* inhabiting this area search for an ecological niche which is able to provide protection as well as a carbon and energy source. Former studies already revealed the coexistence of *A. montana* with arbuscular mycorrhizal fungi (Ryszka et al. 2010), but no reports exist about the discovery of a further inhabitant, *Streptomyces*, in the medicinally used plant of *A. montana*.

As nutrient source the roots of *A. montana* contain carbon hydrates as inulin, glucose, fructose and sucrose. Free amino acids are available as synthesis precursors. Beyond that, a large variety of protective molecules is produced in the roots to keep competitor microorganisms away. Compounds with antibacterial activity are polyacetylenic compounds, essential oils, phenolic ethers and esters. The polyacetylenic molecules additionally provide an antifungal effect. Phenolic acids add an antioxidative facet (Rossetti et al. 1984; Teuscher et al. 2009; Jurkiewicz et al. 2010; Pljevljakušić et al. 2012). Consequently, the benefits for endophytic inhabiting of *A. montana* are apparent.

We found that *Streptomyces* detected in *A. montana* produce several metabolites which can be beneficial for the plant itself. Cycloheximide can be useful to the host plant as an antifungal and antiviral compound (Ji et al. 2010). The antifungal activity could be confirmed by our studies as only cycloheximide-containing extracts 2 and 3, but not cycloheximide-lacking extract 4, showed a

growth-inhibitory effect against *Candida parapsilosis* and *Fusarium verticillioides*. Antifungal effects of cycloheximide against other species of the genera *Candida* and *Fusarium* have already been described (Sharma et al. 2010). Both fungi are present in soil and *Fusarium* is additionally known to be an important plant pathogen (Brown et al. 2006; Nosek et al. 2009). This emphasizes the possible importance of *Streptomyces* producing cycloheximide to protect the host plant against pathogenic fungi. Actiphenol, which has a similar chemical structure, is also known to exhibit an antifungal and antiviral activity as well. Moreover it is discussed to act as a herbicidal agent (Fukuda et al. 2005; Ji et al. 2010). The identified diketopiperazines and the glutarimide antibiotics have been found before in *Streptomyces* (Highet and Prelog 1959; Macherla et al. 2005; Pettit et al. 2006). The role of the different diketopiperazines is multi-layered. The antibacterial activity, which is frequently found in secondary metabolites of *Streptomyces* in order to inhibit competitors (Diminic et al. 2014), is heterogeneously reported. Cyclo-prolyl-valyl exerted no activity against *S. aureus*, *B. subtilis* and *E. coli* (Effendi 2004), but all four diketopiperazines found in our extracts are reported to be active against *Vibrio anguillarum* (Fdhila et al. 2003). Cyclo-leucyl-prolyl is reported to inhibit the growth of *Mycobacterium marinum*, *E. faecalis*, *B. subtilis* and *S. aureus* (Rhee 2002; Effendi 2004; Martins and Carvalho 2007). Notwithstanding, another working group could not show activity of this molecule against *B. subtilis* and *E. coli* (Cabrera et al. 2006). The most active among our four diketopiperazines seems to be cyclo-prolyl-phenylalanyl. A moderate activity against six bacterial strains was already shown for this molecule, among them *E. coli*, *P. aeruginosa*, *S. aureus* and *S. pneumoniae* (Milne et al. 1998). However, we could not find a pronounced antibacterial effect in our *Streptomyces* extracts containing

diketopiperazines except the weak growth reduction of *S. aureus*. This small effect could be explained due to the reported activity of cyclo-prolyl-phenylalanyl (Milne et al. 1998). It is conceivable that the concentration of these moderately antibacterial acting molecules in our extracts is too low to exert more inhibitory activity. Perhaps the low concentration used could also explain why we did not find an activity against *S. aureus* by the diketopiperazine-containing DNPM medium, which was also tested. Beyond that, it is eventually not necessary for the *Streptomyces* strains inhabiting *A. montana* to produce more active antibacterial compounds due to the presence of above-mentioned molecules, who accomplish this function in the *A. montana* root (Jurkiewicz et al. 2010). Other reported activities of the diketopiperazines comprise an antifungal effect of cyclo-leucyl-prolyl and cyclo-prolyl-phenylalanyl, and a phytotoxic effect of cyclo-prolyl-phenylalanyl against certain plants (Holden et al. 1999; Wang et al. 1999; Klose 2006). The most interesting diketopiperazines regarding the interaction with *A. montana* are cyclo-prolyl-valyl and even more cyclo-prolyl-phenylalanyl. In addition to the mentioned antibacterial, antifungal and phytotoxic properties, those molecules seem to promote plant growth. Especially cyclo-prolyl-phenylalanyl is discussed recently to act as a signalling mimic of the plant hormone auxin (Ortiz-Castro et al. 2011; Venturi and Fuqua 2013).

Interestingly, our presumption that those *Streptomyces* could possibly produce sesquiterpene lactones could not be confirmed. Thus, *A. montana*, but not the endophytic *Streptomyces* strains, seems to be the producer of those medicinally effective compounds.

Complementary to the structure identification and examining of the antibacterial and antifungal properties, we tested the impact of the *Streptomyces* extracts on

the viability of Jurkat T cells, a certain sort of T lymphocyte-like cancer cell line. Only those *Streptomyces* extracts which contain cycloheximide and or actiphenol (2, 3, 4 and 5) revealed a concentration dependent reduction in cell viability. Therefore, these extracts can be described as cytotoxic. Both identified glutarimide antibiotics are known to act as cytotoxic substances (Otani et al. 1989; Lakhvich et al. 1993). This reinforces the presumption that those two compounds could be responsible for this activity. Actinobacteria isolated from other members of the Asteraceae family (*Parthenium hysterophorus*, *Ageratum conyzoides*, *Sonchus oleraceus*, *S. asper* and *Hieracium canadense*) were also shown to produce highly cytotoxic substances, but their structures were not elucidated (Tanvir et al. 2014).

In summary, it seems conceivable that those two organisms interact on the one hand to complete their protection against competitors and pathogens and on the other hand to exchange plant growth promoting signals against a carbon and energy source for the microorganism. In our case *A. montana* is highly likely to contribute to this community by offering carbon and synthesis precursors to the *Streptomyces* and protective substances for the defense against competing bacteria and fungi. The *Streptomyces* strains in return could augment the protection against pathogenic fungi by producing cycloheximide and possibly support the growth of *A. montana* by producing cyclo-prolyl-phenylalanyl. All those reasons suggest this symbiosis to take place to facilitate life in this ecological niche of a nutrient-poor and adverse environment. The genomic data of the sequenced strain 3 suggest the huge biosynthetic potential of endophytic bacteria. The number of natural products gene clusters is higher than those of model *Streptomyces* strains such as *S. coelicolor*, *S. lividans* or *S. albus*. Therefore, the exploitation of the endophytic actinobacteria might have a great

impact on drug discovery.

ACKNOWLEDGMENTS

The authors thank Barbara Schuler for her help regarding GC-MS implementation.

COMPLIANCE WITH ETHICAL STANDARDS

Conflict of interest: The authors declare that they have no conflict of interest.

This article does not contain any studies with human participants and animals performed by any of the authors.

REFERENCES

- Ball P (2000) Quinolone generations: natural history or natural selection? *J Antimicrob Chemother* 46:17-24
- Bérdy J (2005) Bioactive Microbial Metabolites. *J Antibiot (Tokyo)* 58:1–26
- Bitzer J (2005) Isolierung und Strukturaufklärung neuer Naturstoffe aus Bakterien und endophytischen Pilzen durch chemisches Screening. Dissertation, University of Göttingen
- Brown DW, Butchko RAE, Proctor RH (2006) *Fusarium* genomic resources: Tools to limit crop diseases and mycotoxin contamination. *Mycopathologia* 162:191–199
- Cabrera GM, Butler M, Rodriguez MA, Godeas A, Haddad R, Eberlin MN (2006) A sorbicillinoid urea from an intertidal *Paecilomyces marquandii*. *J Nat Prod* 69:1806–8. doi: 10.1021/np060315d
- Castillo UF, Browne L, Strobel G, Hess WM, Ezra S, Pacheco G, Ezra D (2007) *Microbial Ecology*. 53:12–19. doi: 10.1007/s00248-006-9129-6
- Coombs JT, Franco CMM (2003) Isolation and Identification of Actinobacteria from Surface-Sterilized Wheat Roots. *Appl Environ Microbiol* 69:5603–5608. doi: 10.1128/AEM.69.9.5603-5608.2003

- Demain AL (1999) Pharmaceutically active secondary metabolites of microorganisms. *Appl Microbiol Biotechnol* 52:455–63
- Diminic J, Starcevic A, Lisfi M, Baranasic D, Gacesa R, Hranueli D, Long PF, Cullum J, Zucko J (2014) Evolutionary concepts in natural products discovery: What *Actinomycetes* have taught us. *J Ind Microbiol Biotechnol* 41:211–7. doi: 10.1007/s10295-013-1337-8
- Effendi H (2004) Isolation and structure elucidation of bioactive secondary metabolites of sponge-derived fungi collected from the Mediterranean sea (Italy) and Bali sea (Indonesia). Dissertation, University of Düsseldorf
- ESCOP (2003) ESCOP Monographs : the scientific foundation for herbal medicinal products, 2nd edn. Thieme, Stuttgart
- Fdhila F, Vázquez V, Sánchez JL, Riguera R (2003) dd-diketopiperazines: antibiotics active against *Vibrio anguillarum* isolated from marine bacteria associated with cultures of *Pecten maximus*. *J Nat Prod* 66:1299–301. doi: 10.1021/np030233e
- Fukuda T, Matsumoto A, Takahashi Y, Tomoda H, Omura S (2005) Phenatic acids A and B, new potentiators of antifungal miconazole activity produced by *Streptomyces* sp. K03-0132. *J Antibiot (Tokyo)* 58:252–9. doi: 10.1038/ja.2005.29
- García-Piñeres AJ, Castro V, Mora G, Schmidt TJ, Strunck E, Pahl HL, Merfort I (2001) Cysteine 38 in p65/NF-kappaB plays a crucial role in DNA binding inhibition by sesquiterpene lactones. *J Biol Chem* 276:39713–20. doi: 10.1074/jbc.M101985200
- Highet RJ, Prelog V (1959) Stoffwechselprodukte von Actinomyceten. 18. Mitteilung. *Actiphenol. Helv Chim Acta* 42:1523–1526. doi: 10.1002/hlca.19590420514
- Holden MT, Ram Chhabra S, de Nys R, et al (1999) Quorum-sensing cross talk: isolation and chemical characterization of cyclic dipeptides from *Pseudomonas aeruginosa* and other gram-negative bacteria. *Mol Microbiol* 33:1254–66
- Ji X, Zhong Z, Xue S, Meng S, He W, Gao R, Li Y, Li Z (2010) Synthesis and antiviral activities of synthetic glutarimide derivatives. *Chem Pharm Bull (Tokyo)* 58:1436–41
- Jiménez-Esquilín AE, Roane TM (2005) Antifungal activities of actinomycete strains associated with high-altitude sagebrush rhizosphere. *J Ind Microbiol Biotechnol* 32:378–81. doi: 10.1007/s10295-005-0007-x
- Jurkiewicz A, Ryszka P, Anielska T, Waligórski P, Białońska D, Góralska K, Tsimilli-Michael M, Turnau K (2010) Optimization of culture conditions of *Arnica montana* L.: effects of mycorrhizal fungi and competing plants. *Mycorrhiza* 20:293–306. doi: 10.1007/s00572-009-0280-z

- Katoh K, Misawa K, Kuma K, Miyata T (2002) MAFFT: a novel method for rapid multiple sequence alignment based on fast Fourier transform. *Nucleic Acids Res* 30:3059–3066. doi: 10.1093/nar/gkf436
- Kearse M, Moir R, Wilson A, et al (2012) Geneious Basic: An integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics* 28:1647–1649. doi: 10.1093/bioinformatics/bts199
- Kieser T, Bibb MJ, Buttner MJ, Chater KF, Hopwood DA (2000) *Practical Streptomyces Genetics*. The John Innes Foundation, Norwich, United Kingdom
- Klose KE (2006) Increased chatter: cyclic dipeptides as molecules of chemical communication in *Vibrio* spp. *J Bacteriol* 188:2025–6. doi: 10.1128/JB.188.6.2025-2026.2006
- Krause L, McHardy AC, Nattkemper TW, Pühler A, Stoye J, Meyer F (2007) GISMO-gene identification using a support vector machine for ORF classification. *Nucleic Acids Res* 35:540–549. doi: 10.1093/nar/gkl1083
- Lakhvich FA, Buravskaya TN, Akhrem AA (1993) Synthesis of actiphenol and its normethyl analogues and 6-hydroxy derivatives. *Chem Nat Compd* 29:526–529.
- Lee K-H, Ibuka T, Wu R-Y, Geissman TA (1977) Structure-antimicrobial activity relationships among the sesquiterpene lactones and related compounds. *Phytochemistry* 16:1177–81
- Lin L, Ge HM, Yan T, Qin YH, Tan RX (2012) Thaxtomin A-deficient endophytic *Streptomyces* sp. enhances plant disease resistance to pathogenic *Streptomyces scabies*. *Planta* 236:1849–61. doi: 10.1007/s00425-012-1741-8
- Loria R, Kers J, Joshi M (2006) Evolution of plant pathogenicity in *Streptomyces*. *Annu Rev Phytopathol* 44:469–87. doi: 10.1146/annurev.phyto.44.032905.091147
- Lyss G, Schmidt T, Merfort I, Pahl H (1997) Helenalin, an anti-inflammatory sesquiterpene lactone from *Arnica*, selectively inhibits transcription factor NF-kappaB. *Biol Chem* 378:951–61
- Macherla VR, Liu J, Bellows C, Teisan S, Nicholson B, Lam KS, Potts BCM (2005) Glaciapyrroles A, B, and C, pyrrolsesquiterpenes from a *Streptomyces* sp. isolated from an Alaskan marine sediment. *J Nat Prod* 68:780–3. doi: 10.1021/np049597c
- Martins MB, Carvalho I (2007) Diketopiperazines: biological activity and synthesis. *Tetrahedron* 63:9923–9932. doi: 10.1016/j.tet.2007.04.105

- Matuschek E, Brown DFJ, Kahlmeter G (2014) Development of the EUCAST disk diffusion antimicrobial susceptibility testing method and its implementation in routine microbiology laboratories. Clin Microbiol Infect 20:O255–66. doi: 10.1111/1469-0691.12373
- Medema MH, Blin K, Cimermancic P, De Jager V, Zakrzewski P, Fischbach MA, Weber T, Takano E, Breitling R (2011) AntiSMASH: Rapid identification, annotation and analysis of secondary metabolite biosynthesis gene clusters in bacterial and fungal genome sequences. Nucleic Acids Res 39:339–346. doi: 10.1093/nar/gkr466
- Merfort I (2010) Arnika – aktueller Stand hinsichtlich Wirksamkeit, Pharmakokinetik und Nebenwirkungen. Zeitschrift für Phyther 31:188–92. doi: 10.1055/s-0030-1262391
- Meyer F, Goesmann A, McHardy AC, et al (2003) GenDB - An open source genome annotation system for prokaryote genomes. Nucleic Acids Res 31:2187–2195. doi: 10.1093/nar/gkg312
- Milne PJ, Hunt a L, Rostoll K, Van Der Walt JJ, Graz CJ (1998) The biological activity of selected cyclic dipeptides. J Pharm Pharmacol 50:1331–7
- Mosmann T (1983) Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. J Immunol Methods 65:55–63.
- Nosek J, Holesova Z, Kosa P, Gacser A, Tomaska L (2009) Biology and genetics of the pathogenic yeast *Candida parapsilosis*. Curr Genet 55:497–509. doi: 10.1007/s00294-009-0268-4
- Ortiz-Castro R, Díaz-Pérez C, Martínez-Trujillo M, del Río RE, Campos-García J, López-Bucio J (2011) Transkingdom signaling based on bacterial cyclodipeptides with auxin activity in plants. Proc Natl Acad Sci U S A 108:7253–8. doi: 10.1073/pnas.1006740108
- Otani T, Sasaki T, Minami Y, Marunaka T, Yu QW (1989) New glutarimide antibiotics, S-632-B1 and B2. I. Taxonomy of producing strain, fermentation and biological properties. J Antibiot (Tokyo) 42:647–53
- Pettit GR, Du J, Pettit RK, Richert LA, Hogan F, Mukku VJR V, Hoard MS (2006) Antineoplastic Agents. 554. The manitoba bacterium *Streptomyces* sp. J Nat Prod 69:804–6. doi: 10.1021/np058087v
- Ph.Eur. (2012) Arnikablüten. In: Eur. Arzneib., 7.3 edn. DAV, Stuttgart, pp 5602–5604
- Pljevljakušić D, Rančić D, Ristić M, Vujisić L, Radanović D, Dajić-Stevanović Z (2012) Rhizome and root yield of the cultivated *Arnica montana* L., chemical composition and histochemical localization of essential oil. Ind Crops Prod 39:177–189. doi: 10.1016/j.indcrop.2012.02.030

- Porras-Alfaro A, Bayman P (2011) Hidden fungi, emergent properties: endophytes and microbiomes. *Annu Rev Phytopathol* 49:291–315. doi: 10.1146/annurev-phyto-080508-081831
- Rhee K-H (2002) Isolation and characterization of *Streptomyces* sp KH-614 producing anti-VRE (vancomycin-resistant enterococci) antibiotics. *J Gen Appl Microbiol* 48:321–7
- Rossetti V, Lombard A, Sancin P, Buffa M, Di Stefano R (1984) Characterization of *Arnica montana* L . Dried Roots. *Int J Crude Drug Res* 22:53–60
- Ruiz B, Chávez A, Forero A, et al (2010) Production of microbial secondary metabolites: regulation by the carbon source. *Crit Rev Microbiol* 36:146–67. doi: 10.3109/10408410903489576
- Ryszka P, Błaszczowski J, Jurkiewicz A, Turnau K (2010) Arbuscular mycorrhiza of *Arnica montana* under field conditions-conventional and molecular studies. *Mycorrhiza* 20:551–7. doi: 10.1007/s00572-010-0302-x
- Sardi P, Saracchi M, Quaroni S, Petrolini B, Borgonovi GE, Merli S (1992) Isolation of endophytic *Streptomyces* strains from surface-sterilized roots. *Appl Environ Microbiol* 58:2691–3
- Sharma N, Thakur M, Chaudhry SC (2010) Monooxovanadium(V) 2-phenylphenoxides: synthesis, characterization, and antimicrobial potential. *J Coord Chem* 63:1071–1079. doi: 10.1080/00958971003710377
- Skwierzynski R, Connors K (1993) Demethylation kinetics of aspartame and L-phenylalanine methyl ester in aqueous solution. *Pharm Res* 10:1174–1180
- Stierle A, Strobel G, Stierle D (1993) Taxol and taxane production by *Taxomyces andreanae*, an endophytic fungus of Pacific yew. *Science* 260:214–216. doi: 10.1126/science.8097061
- Sugier D, Sugier P, Gawlik-Dziki U (2013) Propagation and introduction of *Arnica montana* L. into cultivation: a step to reduce the pressure on endangered and high-valued medicinal plant species. *Sci World J*. doi: 10.1155/2013/414363
- Tamura S, Suzuki A, Aoki Y, Otake N (1964) Isolation of Several Diketopiperazines from Peptone. *Agric Biol Chem* 28:650–652. doi: 10.1271/bbb1961.28.650
- Tanvir R, Sajid I, Hasnain S (2014) Biotechnological potential of endophytic actinomycetes associated with Asteraceae plants: isolation, biodiversity and bioactivities. *Biotechnol Lett* 36:767–73. doi: 10.1007/s10529-013-1430-0
- Teuscher E, Willuhn G, Loew D (2009) *Arnicae flos*. In: Wichtl M (ed) *Teedrogen und Phytopharmaka*, 5th edn. WVG, Stuttgart, pp 91 – 95

- Venturi V, Fuqua C (2013) Chemical signaling between plants and plant-pathogenic bacteria. *Annu Rev Phytopathol* 51:17–37. doi: 10.1146/annurev-phyto-082712-102239
- Wang J-H, Quan C-S, Qi X-H, et al (2010) Determination of diketopiperazines of *Burkholderia cepacia* CF-66 by gas chromatography-mass spectrometry. *Anal Bioanal Chem* 396:1773–9. doi: 10.1007/s00216-009-3379-3
- Wang Y, Mueller UG, Clardy J (1999) Antifungal diketopiperazines from symbiotic fungus of fungus-growing ant *Cyphomyrmex minutus*. *J Chem Ecol* 25:935–941

TABLES

Table 1.

Identified compounds in the respective extracts from the 5 *Streptomyces* strains.

<i>Streptomyces</i> extract no.	cyclo- prolyl- valyl	cyclo- prolyl- isoleucyl	cyclo- prolyl- leucyl	cyclo- prolyl- phenyl- alanyl	cyclo- heximide	actiphenol
1	+	+	+	+	-	-
2	+	+	+	+	+	-
3	+	+	+	+	+	+
4	+	+	+	-	-	+
5	+	+	+	-	+	+

Table 2.

Antifungal activity of *Streptomyces* strains 2, 3, and 4.

	<i>Streptomyces</i> extract no. 2			<i>Streptomyces</i> extract no. 3			<i>Streptomyces</i> extract no. 4			nystatin	methanol
concentration [µg]	10	50	100	10	50	100	10	50	100	75	20 µL
<i>Candida</i> <i>parapsilosis</i>	-	+	+	-	++	++	-	-	-	+	-
<i>Fusarium</i> <i>verticillioides</i>	-	++	++	-	++	++	-	-	-	+	-

- no zone of inhibition, + zone of inhibition lower than the tested 75 µg nystatin, ++ zone of inhibition larger than the tested 75 µg nystatin

FIGURES

Fig. 1

Molecular structures of identified secondary metabolites in *Streptomyces* extracts.

Fig. 2

Cell viability of Jurkat T cells after 24 hours of treatment with *Streptomyces* extracts. Actinomycin D (ActD) represents the positive control and DMSO the negative control. *Streptomyces* extracts were tested in two concentrations, 10 and 50 µg mL⁻¹. Numbers 1 to 5 refer to the respective extracts. Data represent mean ± standard deviation (SD) of two independent experiments. ** $p < 0.01$ and *** $p < 0.001$ vs. DMSO.

Fig. 3

Antibacterial activity against *S. aureus* measured by disc diffusion test. Disk number 1 contained penicillin 1 unit, 2 the solvent control methanol, 3 to 5 extract 4 in the concentrations of 50 (3), 100 (4) and 150 μ g (5), and 6 to 7 nalidixic acid in the concentrations of 15 (6) and 30 μ g (7). A representative susceptibility test is shown.

Fig. 4

Phylogenetic tree of the strains *Streptomyces* sp. *Arnica3* and *S. sp. Arnica5* and related species obtained after 16 sDNA sequencing.