A molecular theory of germinal center B cell selection and division

Graphical abstract

Authors

Michael Meyer-Hermann

Correspondence

mmh@theoretical-biology.de

In brief

Germinal centers are sites in lymphoid tissue, in which high-affinity B cells are selected in an evolutionary process to better fight infections and generate immune memory. Meyer-Hermann provides evidence that the selection and division of B cells is controlled by two different rather than a single molecular signal.

Highlights

- Germinal center B cells use different signals for selection and division
- mTOR, FoxO1, and c-Myc dynamics in B cells can explain conflicting experiments
- Model variants show different light zone passage times for high- and low-affinity B cells
- Clonal diversity may distinguish different model variants

Article

A molecular theory of germinal center B cell selection and division

Michael Meyer-Hermann^{1,2,3,4,5,*}

¹Department of Systems Immunology and Braunschweig Integrated Centre of Systems Biology, Helmholtz Centre for Infection Research, Rebenring 56, Braunschweig 38106, Germany

²Centre for Individualised Infection Medicine (CIIM), Hannover, Germany

³Institute for Biochemistry, Biotechnology and Bioinformatics, Technische Universität Braunschweig, Braunschweig, Germany

⁴Cluster of Excellence RESIST (EXC 2155), Hannover Medical School, Carl-Neuberg-Str. 1, Hannover 30625, Germany

⁵Lead contact

*Correspondence: mmh@theoretical-biology.de

<https://doi.org/10.1016/j.celrep.2021.109552>

SUMMARY

The selection of B cells (BCs) in germinal centers (GCs) is pivotal to the generation of high-affinity antibodies and memory BCs, but it lacks global understanding. Based on the idea of a single Tfh-cell signal that controls BC selection and division, experiments appear contradictory. Here, we use the current knowledge on the molecular pathways of GC BCs to develop a theory of GC BC selection and division based on the dynamics of molecular factors. This theory explains the seemingly contradictory experiments by the separation of signals for BC fate decision from signals controlling the number of BC divisions. Three model variants are proposed and experiments are predicted that allow one to distinguish those. Understanding information processing in molecular BC states is critical for targeted immune interventions, and the proposed theory implies that selection and division can be controlled independently in GC reactions.

INTRODUCTION

Germinal centers (GCs) are specialized environments in lymphoid tissues that give rise to high-affinity antibodies (Victora and Nussenzweig, 2012). The creation of new antibodies that previously did not exist in an organism was well described in many details, including GC morphology (Camacho et al., 1998), with dark and light zones (DZs and LZs), interzonal cell motility (Allen et al., 2007; Schwickert et al., 2007; Hauser et al., 2007; Figge et al., 2008), B cell (BC) mutation (MacLennan, 1994; Oprea et al., 2000), T follicular helper (Tfh) cell-dependent selection and division (Victora et al., 2010; Meyer-Hermann et al., 2012), and clonal diversity (Tas et al., 2016). BCs undergo somatic hypermutation in the GC-DZ, where BC division is dominant, while BC selection is concentrated in the GC-LZ, where native antigen is presented on follicular dendritic cells (FDCs) and cognate interactions with Tfh cells take place.

BC selection is based on two steps (Lindhout et al., 1995): BCs must collect antigen from FDCs and obtain selection signals from Tfh cells. Tfh cells can sense the affinity of the BC receptor (BCR) by the density of antigen-derived peptide-loaded major histocompatibility complex (pMHC) on the BC surface (Meyer-Hermann et al., 2006) and polarize to the BC with the highest pMHC density (Depoil et al., 2005), respective of the highest affinity. When the BC receives a critical amount of Tfh cell signals it is positively selected. BCs failing to do so activate apoptosis. This theory of BC selection based on competition for Tfh cell help was supported by experimental findings (Allen

et al., 2007; Victora et al., 2010) and forms the basis of the present investigation.

The actual signaling of BCs in the course of BC mutation and subsequent selection remains elusive. It was proposed based on *in vitro* experiments that chemical signaling would shape probability distributions of fate decisions, ultimately inducing fate diversity on the BC population level (Duffy et al., 2012). Only recently did molecular BC signaling come into focus in research and point to a critical role of molecules such as c-Myc (Calado et al., 2012; Heinzel et al., 2017; Luo et al., 2018), FoxO1 (Sander et al., 2015; Inoue et al., 2017), and mammalian target of rapamycin (mTOR) (Ersching et al., 2017). All of these factors are involved in BC signaling downstream of BCR and CD40 ligation by Tfh cells, which is critical for BC survival, division, and differentiation, and a synergy of both pathways was identified (Luo et al., 2018; Turner et al., 2018). In addition, a positive feedback loop between CD40 ligation and ICOSL upregulation was described (Liu et al., 2015), giving rise to a non-linear relationship of BC stimulation and downstream signaling, which makes the interpretation of experimental results less intuitive.

One constraint that determines the architecture of signal progression is the observation of a differential number of divisions (DND). Based on *in vivo* observations (Victora et al., 2010) and computer simulation (Meyer-Hermann et al., 2012), it was found that the interaction between Tfh cells and BCs gives rise to BC selection and subsequent differentiation; in addition, the number of subsequent divisions, rather than being a random number or an endogenous BC property, is controlled by these very same

interactions with Tfh cells and ranges between one and six divisions. While this was confirmed in specific experiments (Gitlin et al., 2014; Finklin et al., 2019), it also bears a structural and unresolved limitation of our understanding of the underlying processes: assuming selection was determined by a particular signal X , which is integrated from BC antigen uptake and/or multiple interactions with different Tfh cells, a threshold signal X_{th} would exist at which BC selection and fate decision are initiated. Then, all of the selected BCs would exhibit a similar signal level $X \approx X_{th}$ at the moment of selection. If this very same signal was also responsible for the induced DND in the DZ, then all of the selected BCs would divide an equal number of times, which is not the case (Gitlin et al., 2014). Thus, this selection signal X cannot encode the subsequent DND, the information about which must be encoded in a different signaling molecule Y , which either dynamically controls the threshold X_{th} (Y) or directly controls DND. This article proposes a solution to this structural problem.

The relative role of BCR and Tfh cell signals was disentangled by providing antigen to BCs in a BCR-independent way, taking advantage of the DEC205 receptor (Victora et al., 2010). This receptor is expressed on BCs and allows for antigen-uptake irrespective of the interactions of BCRs with antigen hold on FDCs. BCs that take up antigen via the DEC205 receptor exhibit high antigen-pMHC presentation, prolonged LZ passage times, and a division burst in the DZ. While the division burst can be understood by increased Tfh cell help to high pMHC-presenting BCs, the prolonged LZ passage time was never understood and actually contradicts the general expectation that high-affinity BCs would be selected faster than low-affinity BCs. A realistic theory of BC selection must properly describe GCs permissive of low-affinity BCs (Kuraoka et al., 2016; Silver et al., 2018) still allowing for efficient affinity maturation, diversity of GC clonal dominance (Tas et al., 2016) still allowing for clonal bursts, Tfh cell signal-dependent DND upon selection (Gitlin et al., 2014) without losing the relevance of BCR signaling for selection, and reflect extreme stimulation settings such as antigen uptake via the DEC205 receptor (Victora et al., 2010). A consistent theory that explains these seemingly conflicting GC properties does not exist (Oprea and Perelson, 1997; Zhang and Shakhnovich, 2010; Meyer-Hermann et al., 2012; Wang et al., 2016; De Boer and Perelson, 2017; Meyer-Hermann, 2019).

Here, variants of a hypothetical theory of molecular BC information processing in agreement with all of the aforementioned experimental results are proposed. Our mechanistic understanding of GC-BC selection is improved based on a concept of information processing that separates the signals for selection from those determining the number of subsequent divisions. The three model variants are tested for observable implications in computer simulations and experimental tests are proposed to determine which version of signal separation was realized in nature. As such, the presented simulations must be considered a systematic method for hypothesis testing.

RESULTS

A theory separating selection and division signals

A spatio-temporal GC simulation framework was generated in which each BC, FDC, and Tfh cell is represented as an individual

object (Meyer-Hermann et al., 2012, 2014, 2019; Binder and Meyer-Hermann, 2016). In this framework, BCs divide and mutate in the DZ and interact with FDCs and Tfh cells in the LZ. FDCs harbor an initial amount of antigen, which is taken up by BCs with an affinity-dependent probability, whereby the kinetics of antigen cycling on FDCs is neglected (Arulraj et al., 2021). The subsequent BC fate (i.e., apoptosis or recycling to the DZ phenotype) is determined by signals collected from multiple FDCs and Tfh cells. Note that there is no experimental proof that BCs collect signals from multiple interactions with Tfh cells (Wan et al., 2019). The model variants of the signaling theory described below have in common that they separate the signals for BC fate decision and the number of BC divisions. They detail the mechanisms of BC selection (summarized in Figure 1) and were implemented for each cell object (see STAR Methods and Table 1), inducing a diversity of cell behaviors that depend on the interactions individually experienced by each cell.

In the BCinTime model, BCs of the LZ phenotype receive additional time to search for survival and selection signals by uptake of antigen. Thus, the LZ passage time is determined by the efficiency of antigen uptake from FDCs. This is realized by a phenomenological signal upregulated in LZ-BCs with a rate depending on the amount of uptaken antigen (see Equation 7 and parameters K_R and n_R in Table 1). With every additional antigen, the rate of signal upregulation is reduced. When the signal reaches a threshold value, the fate decision is invoked and the DND is determined by the amount of Tfh cell signals accumulated by then (see Equations 8 and 11 and Table 1).

In the DisseD model (different signals for selection and division), selection and DND are based on the levels of mTOR and c-Myc, respectively (Figure 1). In LZ-BCs, mTOR is upregulated in a BCR-dependent manner (see Equation 10 and parameters K_R and n_R in Table 1). During Tfh cell-BC interactions, BCR and Tfh cell signals synergize to upregulate mTOR more efficiently. When mTOR reaches a threshold expression level, BC fate decision is initiated. During the mTOR-limited LZ passage time, c-Myc is upregulated in a Tfh cell-dependent manner (see Equation 9). Depending on the pMHC level (p), the duration of Tfh cell-BC interactions and the intensity of signaling are modulated (see Table 1, third and fourth block), which affects c-Myc levels. c-Myc upregulation is inhibited by FoxO1. FoxO1 is a transcription factor associated with gene expression during BC development (Dengler et al., 2008) and is dynamically regulated in GC reactions (Dominguez-Sola et al., 2015) where FoxO1 is high in DZ-BCs and low in LZ-BCs (Sander et al., 2015). Thus, c-Myc inhibition by FoxO1 was assumed to be high when BCs enter the LZ. The release of the FoxO1 brake depends on BCR signals (see Equation 4), such that c-Myc upregulation also relies on the synergy of BCR and Tfh cell signals, but with different dynamics. The achieved level of c-Myc determines the subsequent DND in the DZ (see Equation 11 and DND-related parameters in Table 1).

In BCinTime, BCs efficiently collecting or receiving antigen are expected to stay longer in the LZ, as was observed in anti-DEC205-OVA experiments. In contrast, in DisseD, BCs with difficulties in collecting antigen would stay longer in the LZ, as expected for low-affinity cells. The MiXed model aims at reconciling these to seemingly conflicting properties. The mechanisms in DisseD are complemented by a BCinTime element that further limits the LZ

Figure 1. Scheme of the separation of selection and division signals in the 3 models

During their LZ passage time, each simulated BC interacts with FDCs and Tfh cells and collects signals, which determine the fate and time of the fate decision. In BCinTime, a non-specified differentiation signal is upregulated in dependence on the amount of uptaken antigen. This determines the time of fate decision. The amount of Tfh cell signals received by this time determines the DND. In DisseD and MiXed, the signals are made explicit and interact as depicted at the top right in each simulated BC: mTOR is upregulated in dependence on BCR and Tfh cells (CD40). cMyc is upregulated in dependence on Tfh cells (CD40) after BCR signaling-dependent release of the FoxO1 brake. Fate decision is induced when mTOR meets a threshold expression. At this time, BCs with c-Myc, which is subject to degradation, above a threshold survive. The DND is determined by the c-Myc level for surviving BCs. MiXed differs from DisseD by the additional requirement that BCs failing to take up antigen may die before mTOR reaches the fate decision threshold. Implementation in the 3 models is made more concrete in the bottom table. See Table 1 for parameter values.

passage time: LZ-BCs start with an initial finite lifetime, during which they have to manage to catch some antigen from FDCs. Failure to do so implies the irreversible activation of apoptosis. With each successful antigen uptake, the lifetime is prolonged by a fixed time period. Thus, in repeated cycles, each BC must capture new antigen within its limited lifetime. Surviving BCs continue to do so until they reach the mTOR level for fate decision. As in DisseD, c-Myc levels acquired by that time determine the DND.

Basic GC characteristics are similar in all three models

The three models contain a set of parameters specific for the respective signaling strategy (Table 1). These parameters were

adapted to reproduce the generally accepted characteristics of the GC reaction, including reasonable and comparable population kinetics and affinity maturation, a mean DND per DZ around two and in a range between one and six divisions (Meyer-Hermann et al., 2012; Gitlin et al., 2014), a DZ:LZ ratio of two (Victoria et al., 2012), and the right dynamics of this ratio upon anti-DEC205-OVA stimulation (Victoria et al., 2010). Antigen was assumed to be more limited in BCinTime, while the number of Tfh cells was assumed to be more limiting in DisseD and MiXed (Table 1). Parameters for cell motility, somatic hypermutation, affinity maturation, and cell states were kept from previous versions of the model, in which the signaling part was ignored (STAR Methods).

Table 1. Parameter values used in the 3 models in Figure 1

Model name	BCinTime	DisseD	MiXed
Antigen per FDC	3,000	10,000	10,000
Minimum Ag-uptake	1	2	2
LZ-DZ BC-differentiation delay, h	7	6	6
Tfh cell number	300	150	140
LZ passage time given by	Equation 7	Equations 5 and 10	Equations 5 and 10
K_R	4	20	28
n_R	2.4	2	2
Tfh-B interaction time	Gauss	p	p
Min value, h	–	0	0
Max value, h	–	0.3	0.2
K value (p)	–	20	15
Hill coefficient	–	2	2
$T(p)$ (Figure S10G)	1	$T(p)$	$T(p)$
Min value (factor)	–	0	0
Max value (factor)	–	3	4
K value, pMHC (factor = 1)	–	15	15
Hill coefficient	–	2	2
C	Equation 8	Equation 9	Equation 9
$C_{\text{threshold}}$	0.3	0.4	0.2
$\ln(2)/\gamma_C$, h	–	3	3
n_{DND}	1.3	4.0	3.3
C for DND = 2	0.8	0.7	0.82

Parameters were adapted to reproduce experimental constraints (see text and STAR Methods). Equations and symbols refer to the STAR Methods.

The aim of this study was not a quantitative reproduction of available data, which would overinterpret the precision of the data and the simulation results, but a qualitative reproduction of general GC characteristics and the identification of qualitatively different properties depending on the three hypothesized models. In the following, the details of those general characteristics are elaborated. Subsequently, the differences of the three models will be highlighted (summarized in Table 2). The latter will help to design experiments that validate and distinguish the three models.

As GC affinity maturation and GC output is strongly correlated with GC size (Meyer-Hermann et al., 2012), a comparison of different models must rely on reference simulations of similar size (Figure 2A). The thresholds for selection ($C_{\text{threshold}}$) and the parameters controlling mTOR upregulation (K_R and n_R) were chosen correspondingly (Table 1). The GC population is reduced faster for BCinTime because antigen on FDCs is consumed by BCs and access to antigen becomes more difficult (Figures 2B and 2C), which has a double-negative effect on BC survival. It reduces the LZ passage time, which increases the fraction of LZ-BCs dying by apoptosis and reduces Tfh cell-signal levels in surviving BCs, such that they divide less in the DZ. Thus, the faster reduction of the GC population is an inherent result of BCinTime.

Affinity maturation of GC BCs is similar (Figure 2D). While the competition for antigen is critical in BCinTime, this is equilibrated by less competition for Tfh cell help (Figure 2E). The lower affinity of output cells during the first days of the reaction (Figure 2F) points to a more permissive selection in the phase of abundant antigen. The GC accelerates affinity maturation when antigen availability becomes limiting (Figure 2D). However, the ultimate affinity of output cells is kept lower (Figure 2F) for the reduced total number of BCs passing through the GC, which infers less total output per GC in the late reaction (Figure 2G).

In DisseD and MiXed, the information on antigen uptake by the BCR is translated into the speed of release of the FoxO1 brake on c-Myc (Figures 1, S1A, S1B, S1F, and S1G) and stored in the level of pMHC presentation. The latter determines the frequency and the intensity of signals received by Tfh cells, which is reflected in the rate of mTOR and c-Myc upregulation (Figures S1B, S1C, S1G, and S1H). c-Myc levels at the time of fate decision, which is determined by mTOR reaching a threshold value, show a clear separation of deleted and selected BCs. The separation is more pronounced in MiXed (Figures S1D, S1E, S1I, and S1J). The range of c-Myc levels in the subset of selected BCs encodes the DND.

The DND parameters were chosen to generate a mean value ~ 2 in a range of 1–6 (Figures 2H and 2I), which is consistent with the results of Meyer-Hermann et al. (2012) and Gitlin et al. (2014). A mean number of 2 divisions per round is supported by the DZ:LZ ratio of 2, which is a needed starting point for the anti-DEC205-OVA experiments (see below). In BCinTime, the DZ:LZ ratio needed to be adjusted by a longer CC-CB differentiation delay (Table 1). As a result of less-efficient antigen uptake, low-affinity BCs divide less than high-affinity BCs in all models. While for DisseD and MiXed, DND drops in the phase of strong competition for Tfh cell help, in BCinTime, DND increases in the same phase and drops afterward (Figure 2H). This reflects the critical role of antigen availability in BCinTime.

anti-DEC205-OVA experiments explained *in silico*

It is possible to provide antigen to BCs expressing the DEC205 receptor via the injection of anti-DEC205-OVA. The BCs then internalize, process, and present the antigen as pMHC on their surface, just as they do with antigen taken up via the BCR but without activating the signaling machinery downstream of the BCR and without the need to actually interact with FDCs to catch antigen. Anti-DEC205-OVA injection *in vivo* induced an accumulation of BCs in the LZ for 12 h before a burst of BC division in the DZ, as reflected in the dynamics of the DZ:LZ ratio (Victoria et al., 2010). Previous GC models, such as the LEDA model (Meyer-Hermann et al., 2012), were able to explain the DZ burst, but failed to explain the LZ accumulation without additional assumptions, such as increased interaction times with Tfh cell or equivalent, which were not supported by experimental data. Next, the consistency of the three selection models with LZ accumulation and DZ burst was tested.

Injection of anti-DEC205-OVA *in silico* leads to an accumulation of BCs in the LZ and to a subsequent proliferation burst in the DZ that is consistent with the experimental data in all of the models (Figures 3A–3C). For DisseD and MiXed, strong non-linearities in the dependence of DND on c-Myc levels (see n_{DND} in

Table 2. Qualitative experimental observations and how they are reflected in the 3 models

Observation/reference	BCinTime	DisseD	MiXed	Figure
GC size (Liu et al., 1991; Hollowood and Macartney, 1992)	reasonable GC size and population kinetics			2A
GC shutdown (Zhang et al., 2013)	fast	slow	slow	2A
Robustness against antigen concentration (Hannum et al., 2000)	low	high	high	2B, S4A, S4B, S4D, S4E
Affinity maturation (Victora et al., 2010)	well reproduced	well reproduced	well reproduced	2D
DZ:LZ ratio is 2 (Victora et al., 2010)	well reproduced	well reproduced	well reproduced	3A–3C
Competition for Tfh (prediction)	low	high	high	2E
Variation of DND between 1 and 6 (Meyer-Hermann et al., 2012; Gitlin et al., 2014)	assumed	assumed	assumed	2H, 2I, 3G, S2I, S4B, S4E
Mean DND of ≈ 2 (Meyer-Hermann et al., 2012)	2.23 ± 0.03	2.16 ± 0.03	2.0 ± 0.04	2H, 2I, 3G, S2I, S4B, S4E
Asymmetry of DZ to LZ and LZ to DZ transmigration (Victora et al., 2010)	well reproduced	well reproduced	well reproduced	4A
LZ passage time of high-affinity BCs (prediction)	longer	shorter	shorter	4B, S4C, S4F, S5A
LZ passage time of deleted BCs (prediction)	shorter	longer	shorter	4C, S5B
Number of Tfh cell contacts per LZ passage in deleted BCs (prediction)	less	more	less	4D, S5C
LZ accumulation and DZ burst in anti-DEC205-OVA experiments (Victora et al., 2010)	well reproduced	well reproduced	well reproduced	3A–3C, S2A–S2C
Stop of affinity maturation in anti-DEC205-OVA experiments (Victora et al., 2010)	well reproduced	well reproduced	well reproduced	S3
mTOR-overexpressing BCs have competitive disadvantage (Ersching et al., 2017)	qualitatively reproduced	qualitatively reproduced	qualitatively reproduced	S4G, S4H
Clonal diversity (Tas et al., 2016)	qualitatively reproduced	qualitatively reproduced	qualitatively reproduced	5A–5C
Existence of monoclonal and multiclonal GCs (Tas et al., 2016)	yes, frequent monoclonal	yes	yes, frequent multiclonal	5D, 5E, S6A–S6D
Clonal dominance day 17 of the GC reaction (prediction)	higher	lower	intermediate	5G

Table 1) were not sufficient to receive an appropriate burst. Thus, a pMHC-dependent Tfh cell-BC interaction time and Tfh cell signaling ($T(p)$ in Table 1 and Figure S10G) were introduced. For simplicity, the Hill coefficient of both mechanisms was fixed to 2. All three non-linearities together were able to generate DZ bursts in this setting. The degree of non-linearities can be shifted between the three assumed mechanisms.

LZ accumulation is robust against changes of the amount of DEC205⁺ BCs in BCinTime, but not in DisseD and MiXed, for which the LZ accumulation phase is extended and DZ expansion is reduced with more DEC205⁺ BCs (Figures S2A–S2C), because stronger competition for Tfh cells leads to slower upregulation of mTOR and c-Myc. The DZ burst relies on dynamics in DND (Figures 3G and S2I), with lower peak DND in BCinTime. While in DisseD and BCinTime, the GC relaxes to normal within 3 days, the advantage of DEC205⁺ BCs persists for longer times in MiXed (Figure 3C). Similar to the *in vivo* case, the DEC205⁺ population exhibits only small modulations by the *in silico* injection. The fewer the DEC205⁺ BCs in the system, the weaker the

perturbation of the GC and the effect on DEC205⁺ BCs (Figure S2A–S2C).

In BCinTime, LZ accumulation is a natural result of the principle that BCs receive additional Tfh cell search time with every up-taken antigen. The amount of pMHC is increased by the *in silico* injection, and so is the LZ passage time by definition (Figure 3D, rectangle). In the case of DisseD and MiXed, LZ accumulation can only be understood by the analysis of mTOR, FoxO1, and c-Myc dynamics. The injection of anti-DEC205-OVA infers high pMHC, while signals downstream of BCR are weak. Low BCR signaling has an impact on FoxO1 and mTOR (Equations 4 and 10), thus inducing a slower upregulation of mTOR despite high pMHC levels (Figures 3E and 3F, rectangles, and S2D), which explains the LZ accumulation. Low BCR also delays the release of the FoxO1 brake on c-Myc (Figure S2E). As the DEC205⁺ BCs continue to collect antigen from FDCs, they finally receive some BCR signals. In the model, FoxO1 was assumed to be more sensitive to low BCR signals than mTOR; thus, the FoxO1 brake is released first, allowing for c-Myc upregulation,

Figure 2. General GC characteristics of the three models

(A) Population kinetics, (B) antigen collection and (C) consumption, (D) mean affinity of GC-BCs, (E) competition for Tfh cells, (F) mean affinity and (G) number of all output cells produced per GC, (H) mean, and (I) distribution of the number of divisions attributed to selected BCs. DisseD (black), BCinTime (red), and Mixed (blue). Mean and standard deviation (SD) over 100 GC simulations (A–G and I) and over all GC-BCs in a single representative simulation (H).

while mTOR upregulation is kept slow (Figures S2D and S2F). Because of the longer LZ passage time, together with more intense and longer Tfh cell-B cell interactions, the BCs have the chance to interact with many Tfh cells, such that BCs exhibit higher c-Myc levels at the time of selection (Figures 3H and 3I, rectangles, and S2G and S2H, green lines), subsequently leading to the DZ burst (Figures 3G and S2I, gray and blue).

The range of c-Myc levels at selection increased between days 3 and 9 of the GC reaction (Figures 3H and 3I, outside rectangles). This reflects increasing antigen limitation with time together with affinity maturation in the DEC205⁺ BC subpopulation and holds in normal GC simulations (without anti-DEC205-OVA injection), and allows for the diversity of DNDs of selected BCs (Figure 2I).

Figure 3. Providing antigen via the DEC205 receptor

Following the experimental protocol (Victora et al., 2010), simulations were started from high-affinity founder clones at a distance of 2 mutations to the optimal clone and with 20% DEC205⁺ BCs (80% negative). Anti-DEC205-OVA was injected *in silico* at day 5 post-GC onset, which was assumed to increase pMHC presentation to the equivalent of 100 antigen uptake events in all unselected DEC205⁺ BCs in the LZ in a time window of 24 h. Compared to Figure 2B, this corresponds to a 5-fold higher pMHC presentation, as seen in Victora et al. (2010).

(A–C) DZ:LZ ratio in the 3 models BCinTime (A), DisseD (B), MiXed (C), and comparison to experimental data (symbols) from Victora et al. (2010).

(D–F, H, and I) LZ passage time (D–F) and c-Myc (H and I) in dependence on pMHC in BCinTime (D), DisseD (E and H), and MiXed (F and I). Each dot represents 1 cell from a single simulation. Color codes time (see inset). Green and blue dots in rectangles are anti-DEC205-OVA-targeted DEC205⁺ BCs.

(G) Distribution of the DND.

Mean and SD of 100 simulations for each model in (A–C and G).

Consistent with the *in vivo* results (Figure 7G in Victora et al., 2010), affinity maturation is stopped by anti-DEC205-OVA injection *in silico* (Figure S3) because BCs show high pMHC to Tfh cells irrespective of the actual quality of the BCR. The production of output cells in response to anti-DEC205-OVA injections was markedly enhanced in all three models (Figure S3G), but not to the extent seen *in vivo*, suggesting a polarization of BC fate decisions toward the plasma cell phenotype in the case of high pMHC presentation.

LZ passage times differ in the three models

Migration between the DZ and LZ and back is a characteristic property of GC reactions and was quantified by the photoactivation of cells in each of the GC zones (Victora et al., 2010). This revealed a strong asymmetry between both directions of transmigration. The new parameters associated with molecular signaling mechanisms do influence the LZ and DZ passage times and the number of divisions in the DZ. They, therefore, also do affect transmigration rates. This readout was not fine-tuned by parameter variation, but was still reproduced by all three models (Figure 4A), with a tendency to underestimate DZ to LZ migration, thus providing an independent validation of the models.

The LZ passage time (i.e., the time a BC takes from differentiation to the LZ phenotype until initiation of back differentiation to the DZ phenotype) differs in the three models. While in BCinTime, the collection of more pMHC leads to longer times in the LZ (Figure 3D, ignoring the rectangle); the contrary is the case for DisseD and MiXed (Figures 3E and 3F, ignoring the rectangle). One may intuit that high-affinity BCs are more efficient in receiving selection signals and thus would have a shorter LZ passage time. In DisseD (Figure 4B, black lines), low-affinity BCs take 8 h, while high-affinity BCs manage to pass through the LZ state in 5 h. Later, when antigen becomes limiting, both times become longer. In BCinTime (red lines), high-affinity BCs take longer to pass through the LZ. When antigen becomes limiting, the LZ passage times become shorter. In MiXed (blue lines), the high- and low-affinity BCs behave similar to DisseD because high-affinity cells reach the mTOR threshold earlier. The increase in LZ passage times when antigen becomes sparse is less pronounced. The LZ passage times of BCs with low and high affinity are robust against changes in the total amount of antigen used for *in silico* immunization (Figures S4C and S4F). The Pearson correlation coefficients of the LZ passage time with BC affinity at day 5 post-GC onset were -0.76 (95% confidence interval [CI] -0.79 to -0.73) for DisseD, -0.73 (95% CI -0.77 to -0.69) for MiXed, and $+0.78$ (95% CI $+0.76$ to $+0.81$) for BCinTime. The correlation becomes weaker over the time of the reaction, but the difference in sign is maintained throughout (not shown).

Next, the LZ passage time of deleted and selected BCs was compared (Figure 4C). Deleted BCs exhibit a short LZ passage time in the two models with antigen uptake-dependent LZ time (i.e., in BCinTime and MiXed), and a long passage time in DisseD. Because of weak BCR signaling, mTOR is not upregulated quickly in the BCs (see Equation 10), such that the fate decision threshold is reached at rather late times or not reached at all if the BCs hit an imposed maximum LZ passage time. Thus, the LZ passage time of the deleted BCs is diametrically different in

DisseD. Consequently, the number of interactions with Tfh cells per deleted BC, which is in the range of 10–20 per LZ passage for selected BCs, is reduced to a few contacts in BCinTime and MiXed, while it is increased to up to 25 but inefficient contacts in DisseD (Figure 4D).

The BCinTime model is more sensitive to antigen titration

In vivo GC kinetics turned out rather robust against variations of the amount of antigen or of immune-complex deposition on FDCs in response to 4-hydroxy-3-nitrophenylacetyl (NP)-hapten carrier constructs in mice (Hannum et al., 2000). Reducing the antigen by a factor of five *in silico* reduced the peak GC population to one-third in all of the models (Figure S4A). Increasing the antigen by a factor of five retarded GC shut down (Figure S4D). This also induced a 2- to 3-fold peak population in BCinTime, but only a small increase in DisseD and MiXed (Figure S4D). Thus, the GC reaction in BCinTime is more sensitive to the total amount of antigen and less sensitive to Tfh cells (see also Figures 2B and 2E). Despite substantial changes in the total population, DND was independent of antigen in all of the models (Figures S4B and S4E) and is a robust GC property (Gitlin et al., 2014).

All models are consistent with mTOR overexpression experiments

mTOR overexpressing BCs, when coexisting with normal BCs in GCs, have a competitive disadvantage, exhibit a larger DZ:LZ ratio, and hardly affinity mature, as was observed in C57BL/6 mice immunized with NP-OVA (Ersching et al., 2017). This was observed by mixing BCs that express Tsc1 normally with BCs deficient in Tsc1. Tsc1 is an inhibitor of mTOR, and the Tsc1-deficient BCs express more mTOR. In DisseD and MiXed, a larger starting level of mTOR was imposed in BCs at LZ entry (Figures S4G and S4H). As *in vivo*, the DZ:LZ ratio was enhanced to 3–5 in the Tsc1-deficient BCs ~days 4–5 post-GC onset (day 7 post-immunization). These BCs disappeared faster from the GC than *in vivo*. Affinity maturation was inhibited in Tsc1-deficient BCs similar to the *in vivo* case (data not shown). In BCinTime, mTOR is replaced by an unspecified differentiation signal. Imposing a larger starting value of this signal is a phenomenological representation of Tsc1 deficiency and induces the same phenotype as described above (Figures S4G and S4H; red lines). Release of mTOR inhibition *in silico* induced a GC phenotype that is consistent with *in vivo* observations. This supports the credibility of the proposed signaling models.

Clonal dominance gradually differs in the models

It was shown before that GCs in response to chicken gamma globulin or NP-ovalbumin exhibit a great diversity of how strong founder BCs dominate a reaction (Tas et al., 2016). Either a single GC founder clone bursts and then dominates the GC or many founder clones coexist during the whole GC reaction. This was revealed by staining GC BCs with 1 of 10 random colors (confetti mice), which is kept in all progeny, and subsequently observing the dominance of the color in the course of the reaction. The same staining procedure was applied *in silico*. As in experiments, a large diversity of color dominance is found (Figures 5A–5C) in all three models. The diversity of color dominance rises

Figure 4. Transzone migration and LZ passage time for high- and low-affinity B cells

(A) GC simulations were mounted with high-affinity founder BCs at a distance of 2 mutations to the optimal clone, which reflects the experimental setup based on B1-8^{hi} BCs with high specificity for the hapten 4-hydroxy-3-nitrophenylacetyl (NP) (Victora et al., 2010). BCs in the DZ (full lines) or in the LZ (dashed lines) were marked *in silico* and tracked. The fraction of marked BCs found in the opposite zone was measured over time and is shown for the 3 models (colors). Mean and SD of 100 simulations. Experimental data reproduced from Victora et al. (2010).

(B) The LZ passage time (with random founder BCs) is distinguished for high- (full lines) and low- (dashed lines) affinity BCs in the 3 models. High affinity is defined as a binding probability of >0.6 (Equation 1).

(C) Same as (B) for selected and deleted BCs.

(D) Number of interactions with Tfh cells per BC in 1 LZ passage period.

In (B–D), the mean of all BCs of a representative GC simulation for each model is shown (for SDs, see Figure S5). See Figures S4C and S4F for (B) with different levels of antigen.

faster in DisseD and MiXed, and the increase in the baseline dominance observed *in vivo* is only recapitulated in BCinTime and MiXed.

While it was shown previously that color and clonal dominance correlate (Meyer-Hermann et al., 2018), simulations offer the possibility of following the abundance of every clone, which corresponds to sequencing every GC cell at multiple time points of the GC reaction. Both clonal bursts leading to monoclonal GCs

and long-term coexistence of many clones in a single GC were observed *in silico* (Figures 5D and 5E), irrespective of the model (Figures S6A–S6D). This was not associated with a different level of affinity maturation (Figure S7). The clonal dominance distribution over many GCs unraveled gradual differences between the models (Figures 5F, 5G, and S6E–S6J): Clonal dominance starts low and increases to levels ~30% dominance by day 5 in all of the models. Later on, only a few monoclonal GCs evolved in

Figure 5. Clonal dominance

(A–C) Experiment with confetti mice, in which 1 of 10 colors is activated by tamoxifen. Data (black open squares) reproduced from [Tas et al. \(2016\)](#). Following the experimental protocol, 1 of 10 colors is attributed to GC BCs with given probabilities (Table 1 of [Meyer-Hermann et al., 2018](#)). Colors are inherited by daughter cells; the dominance of colors is monitored and reported at different GC time points for DisseD (A), BCinTime (B), and MiXed (C) (gray open squares).

(D and E) Muller diagram showing the relative abundance of founder clones (colors) over the course of the GC reaction (time resolution, 1 h). Two extreme cases are shown for MiXed. See [Figures S6A–S6D](#) for the other models.

(F and G) Each founder clone and its progeny are tracked *in silico*. The largest fraction of BCs stemming from 1 clone is monitored. The number of GCs among 1,000 GCs with a most dominant clone of particular dominance (horizontal axes) is depicted at days 5 and 17 post-GC onset (p.o.). See more time points in [Figure S6](#).

DisseD, and the dominance level was kept in the same range. In MiXed, the dominance level further increased, and a small subset of GCs developed into monoclonal GCs. In BCinTime, monoclonal GCs appeared in 20% of the GCs. The other 80% were distributed over a wide range of dominance, with an overall increased dominance level.

This observation reflects the dynamics of the limiting factor in the three models. In BCinTime, antigen is the most limiting, and reduced antigen availability increases selection pressure over the GC reaction. In contrast, in DisseD and MiXed, Tfh cell help is the most limiting. Tfh cell help does not become more limiting over the GC reaction, and in the phase of reduced B cell population, competition for Tfh cell help may even be reduced. This allows for more different B cell types to coexist over a longer period in the GC reaction. Consequently, the three models can

be distinguished in experiments by the frequency of monoclonal GCs. Note that the absolute frequencies of monoclonal GCs depend on the quality of the GC founder BCs ([Figure S8](#)).

DISCUSSION

Separation of selection and activation signals

Three variants of a theory of GC BC selection were proposed on the level of intracellular antigen processing and molecular signaling. Previous GC selection models relied on a single global Tfh cell-signaling level that controls BC selection and subsequent division. These models did not allow the capture of all of the aforementioned experimental results, which is now achieved by all three models presented here (see the summary in [Table 2](#)). In particular, the physiologically extreme anti-DEC205-OVA

experiments (Victoria et al., 2010) now appear as a natural result of the selection mechanisms at work. The contradiction between the assumed selection threshold and the DND of the recycled BCs was resolved in all three models by the separation of selection and activation signals. The theory of separation of those signals complements the known spatial separation of division and selection into DZ and LZ and solves a long-standing inherent problem of GC selection models.

In DisseD and MiXed, the time point of selection of BCs is determined by BCR- and Tfh cell-dependent mTOR upregulation. mTOR upregulation was found in experiments, as indicated by downstream phosphorylation of S6 (Luo et al., 2018). This theory is consistent with the finding that mTOR is necessary for the recycling of GC BCs, but the blocking of mTOR after selection does not affect the number of divisions (Ersching et al., 2017). The exact rule of how to integrate BCR and Tfh cell signals is not known. In the model, mTOR can be upregulated by both signals, such that sufficiently strong BCR signals alone may induce mTOR. As BCR signals are assumed to be a prerequisite of mTOR upregulation by Tfh cell signals (Equation 10), BCR and Tfh cell signals synergize with each other.

According to the models, the number of divisions is determined by Tfh cell-dependent c-Myc levels acquired in the mTOR-determined LZ passage time. This allows for a DND as observed *in vivo* (Gitlin et al., 2014). c-Myc is a powerful driver of anabolic metabolism and cell growth (Wilhelm et al., 2016), which was associated with the selection and recycling of GC BCs (Dominguez-Sola et al., 2012), but without the distinction of selection versus DND. c-Myc is essential for GC maintenance (Calado et al., 2012) and acts as a timer for BC division (Heinzel et al., 2017; Finkin et al., 2019). *In vitro*, there is further evidence that chemical signal like c-Myc would determine the fate distribution of BCs (Duffy et al., 2012). The assumed fast c-Myc decay increases the selection pressure because less frequent encounters of Tfh cells induce the loss of c-Myc in between two interactions with Tfh cells. Thus, BCs with a high frequency of productive BC-Tfh cell interactions (i.e., those with high pMHC) preferentially manage to achieve higher c-Myc scores. Without c-Myc decay, low-affinity BCs may acquire higher c-Myc scores because of their longer LZ passage time.

In the simulation, c-Myc upregulation is inhibited by FoxO1 (see $(1 - H_C(F))$ in Equation 9). The FoxO1 brake on c-Myc was released in dependence on BCR signals, as suggested by data (Luo et al., 2018). The activation of Akt and phosphatidylinositol 3-kinase (PI3K) downstream of BCR (Tzivion et al., 2011) translocate FoxO1 to the cytosol (Burgering, 2008), which leads to FoxO1 inactivation (Brunet et al., 1999). This BCR-dependent mechanism further increases the competitive advantage of high-affinity BCs. Release of the FoxO1 brake opens the path for the Tfh cell-dependent upregulation of c-Myc via CD40 ligation and the NF- κ B pathway (Luo et al., 2018). Therefore, BCR and Tfh cell signals synergize to upregulate c-Myc.

In BCinTime, the separation of selection and division signals is realized in a phenomenological language. The LZ passage time, and consequently the selection signals, are now strongly coupled to antigen uptake rather than to Tfh cell signals. This infers a higher sensitivity to variations of antigen concentration (see below).

FoxO1 is more sensitive to BCR signals than mTOR

The assumption that c-Myc upregulation via release of the FoxO1 brake relies on BCR signals in DisseD may be considered in contradiction to the anti-DEC205-OVA experiments (Victoria et al., 2010), which showed a BCR signaling-independent high DND. However, the prolonged LZ phase still allows for an accumulation of c-Myc signals *in silico* despite the inhibition of Tfh cell signals in BCs with low BCR signals. Also, in reality, despite the removal of Syk signaling downstream of BCR, the partial upregulation of c-Myc was still found (see Figure 7 in Luo et al., 2018). In the model, anti-DEC205-OVA-targeted BCs continue interacting with FDCs, such they receive a minimum of BCR signals. c-Myc upregulation becomes possible provided FoxO1 inhibition is more sensitive to BCR signals than mTOR (as guaranteed by the condition $K_F < K_R$; see STAR Methods). Then, low BCR signals prolong the LZ phase by slow mTOR upregulation, but allow for c-Myc signals because of the moderate release of the FoxO1 brake.

The models differ in the GC clonal composition

The current experimental data situation does not allow us to determine a clear preference for any of the three models. The diversity of clonal dominance in different GCs is found in all of the models (Figures 5A–5C). This relies on a higher degree of non-linearities in DisseD and MiXed (see Table 1) because of more subtle differences in the achieved c-Myc levels, which still must induce a DND. The degree of non-linearities was in particular determined by the strong division burst observed in response to the anti-DEC205-OVA experiment (Victoria et al., 2010). Comparison of *in vivo* and *in silico* experiments using the Brainbow allele, which allows the tracking of BC clones and their progeny through the GC reaction (Tas et al., 2016), favors BCinTime (Figures 5A–5C). However, the frequency of mono-colored GCs depends on the choice of the set of founder BCs. Furthermore, the number of GCs investigated *in vivo* is not sufficient to draw a conclusion. Next-generation sequencing of all GC BCs at late time points of the reaction would allow us to better distinguish the models (Figures 5F and 5G). The simulations predict higher clonal dominance and a large fraction of single-clone-dominated GCs in BCinTime.

GCs are more permissive to B cells with low BCR signals in DisseD

In DisseD, the integration of BCR and CD40 signals for the upregulation of c-Myc (Equation 9) includes dynamics from a positive feedback loop between ICOSL and CD40 (Liu et al., 2015) and a BCR-dependent release of the FoxO1 brake. In reality, BCR signals release the brakes on Tfh cell-mediated signals via the nuclear factor κ B (NF- κ B) pathway (Luo et al., 2018), which is phenomenologically included in the FoxO1 brake. A direct impact of BCR signals on c-Myc upregulation was not considered. BCR signals must be strong enough (as may be monitored by Syk phosphorylation in reality), such that Tfh cell signals can induce a relevant effect on c-Myc upregulation. Low-affinity BCs, i.e., inefficient antigen collectors, may still be selected based on slow but persistent mTOR activity, but receive poor division signals (i.e., low DND). This is consistent with data showing low-affinity survivors in GCs (Kuraoka et al., 2016) or

the persistence of unrelated BCs in the GC reaction (Silver et al., 2018). More generally speaking, GCs in DisseD are more permissive than in BCinTime. This implies that the diversity of BCs generated by the GC reactions tends to be larger in DisseD, being associated with a larger flexibility of the memory BC repertoire to mutations of the pathogen.

High- and low-affinity B cells differ in their LZ passage times

A clear-cut distinction of the models is possible by a measurement of the LZ passage times of high- and low-affinity BCs (Figure 4B), which turned out to be a robust property against the variation of antigen levels (Figures S4C and S4F). In BCinTime, high-affinity BCs stay longer than low-affinity BCs in the LZ. This is inverted in DisseD and MiXed, in which the LZ passage time is determined by mTOR in dependence on both BCR and Tfh cell signals. In high-affinity BCs, higher BCR activation induces faster mTOR upregulation (Equation 10), productive BC-Tfh cell contacts become more frequent, and each contact provides more intense signals to the BC (see $H_L(t)T(p)$ in Equation 10). This reflects synapse intensity and a positive feedback loop involving ICOSL upregulation in BCs (Liu et al., 2015), together, inducing faster selection. In MiXed, BCs must collect antigen for survival (as in BCinTime), but the ultimate LZ passage time is determined by mTOR (as in DisseD). The drastic increase in LZ passage times found in DisseD in the late reaction was not observed in MiXed.

A possible setup for a decisive experiment would use two BC lines with different affinity for the antigen in an activation-induced cytidine deaminase-knockout (AID-KO) system with suppressed mutations. Even with ongoing mutations, a setup with B1-8^{hi} cells together with OVA immunization would allow the monitoring of high- versus low-affinity BCs (Stoler-Barak et al., 2019). Tracking those cells through the LZ would reveal the LZ passage time of both BC types. In a measurement at approximately day 10 post-immunization, a 2-photon setting lasting for 10 h would be needed. The relative LZ passage times of high- and low-affinity BCs are more informative than the absolute times. Relative times were kept throughout the GC reaction, while absolute times varied and were more sensitive to different experimental setups. In BCinTime, LZ passage times are higher at the beginning of the reaction, when antigen is high, and decrease later on. In DisseD and MiXed, the time course is inverted. Immunization with large antigen doses or with an antigen for which high-affinity BCs preexist are predicted to induce different time courses of LZ passage times over the GC reaction.

If it were possible to monitor the time spent in the LZ of BCs with the fate to die, this time would be in the range of 10 h in DisseD and only of 3 h in BCinTime and MiXed.

Mechanisms controlling the LZ passage time

In BCinTime and MiXed, each antigen uptake event is associated with a prolongation of the time period of collecting signals from FDCs and Tfh cells in the LZ. Mechanistically, the determination of the LZ time must be independent of BCR signals because providing the antigen via the DEC205 receptor also induces a prolongation of the LZ passage time *in vivo* (Victoria et al., 2010), as recapitulated in BCinTime (Figure 3A). The prolongation of the LZ time cannot rely on the antigen-processing appa-

ratus either, because providing an irrelevant antigen via the DEC205 receptor does not induce a longer LZ passage time (Victoria et al., 2010). A possible scenario is implemented in MiXed, in which the BC lifetime is prolonged with antigen uptake, reflecting the old idea of the suppression of pre-activated apoptosis by antigen uptake (MacLennan, 1994). The inhibition of centrocyte apoptosis is believed to depend on the interaction with FDCs and Tfh cells (Lindhout et al., 1995; Brandtzaeg, 1996; Tew et al., 1997; Hollmann and Gerdes, 1999; Hur et al., 2000; van Eijk et al., 2001). In the present context, only the interaction with Tfh cells can be responsible for suppressing apoptosis because the prolongation of the LZ phase happens independently of interactions with FDCs. Alternatively, rather than targeting apoptosis, Tfh cells may prevent the acquisition of the DZ phenotype in highly pMHC-loaded BCs by the suppression of FoxO1 via the activation of Akt (Luo et al., 2018). The activation of Akt would be stronger in high-pMHC BCs, thus enforcing higher levels of mTOR to suppress Akt (Ersching et al., 2017) and allowing for FoxO1 upregulation; this delays the recycling and acquisition of the DZ phenotype.

Antigen doses are more relevant in BCinTime

In other models, the size of GCs was found to be dependent on the amount of antigen (Garg et al., 2019). The persistence of antigen availability was shown to increase the amount of resulting antibodies in computer simulations and C57BL/6 mice (Tam et al., 2016). Similarly, slow versus fast antigen delivery was found to be relevant to the GC response in rhesus macaques (Cirelli et al., 2020). In the three models presented here, antigen titration revealed a robustness of the number of divisions per selection round despite rather different population kinetics. BCinTime simulations were the most sensitive to the amount of antigen presented on FDCs, while MiXed and DisseD relied more on Tfh cell help. In view of the observed robustness of the GC reaction against different antigen doses (Hannum et al., 2000), this may be considered an argument against BCinTime. In the context of vaccination based on individual shots of antigen, the resulting GC response would depend less on the vaccine dose in DisseD and MiXed.

Speculation on adaptive control by Tfh cells

In view of the robustness of GC reactions kinetics to changes in the levels of immune complex deposition on FDCs (Hannum et al., 2000), one may speculate whether Tfh cell help is adaptive to the overall level of pMHC levels detected on BCs, thus allowing for a dynamic regulation of the strength of competition for Tfh cell help. In such a model, Tfh cell signaling intensity would be gauged by previously seen pMHC densities, thereby allowing for the proper support of BCs in a framework of low antigen availability.

Evolutionary reasons for signal separation

With computer simulations, the separation of signals for the speed of selection in the LZ and for the number of divisions in the DZ was identified as a concept of information processing in GCs that reconciles seemingly contradicting experimental data. The separation of signals is a well-known concept (e.g., in digital audio signals) that enables independent modulation of

different parts of information. Evolutionarily, this allows for a larger diversity of BCs surviving GC selection, but is limited by their low division potential—thus keeping the focus on the need for high-affinity antibodies in the acute situation. The survival of low-affinity cells is reflected in the diversity of clonal dominance in GCs and is critical to the flexibility of the immune system facing fast-mutating pathogens.

Evolutionary differences of the three model variants of the theory

From an evolutionary perspective, DisseD and MiXed appear advantageous. High-affinity cells are selected faster, which may be important in view of a time-critical fight against an infection. Robustness against the variation of the amount of antigen is higher, which makes the immune system less vulnerable to immune-evasion strategies of slowly expanding pathogens. Fewer GCs are monoclonal, inferring a better preparation for future infections with mutated pathogens by a diverse memory repertoire. The presented computer simulations predict testable readouts that allow the distinguishing of the three proposed models, and by that, to shed light on the mechanisms at work in real GCs and paving the way for improved vaccination protocols. The author hopes to stimulate further experimentation along these lines.

Limitations of study

The present investigation relies mostly on data generated with the hapten-NP-ovalbumin immunization protocol (Rajewsky et al., 1969; Katz et al., 1970). It is known that GC dynamics depends on the choice of hapten versus protein antigen (Zinkernagel, 1997). Also, the model does not consider differences in BC division in dependence on antigen valency (Kato et al., 2020). Thus, the presented investigation of different hypotheses on BC selection and division exclude any application to specific differences in the type of antigen. Furthermore, the representation of antigen-antibody interactions in a phenomenological shape space excludes a deeper analysis of antibody cross-reactivity to antigens, as considered in other models (Wang et al., 2015; Robert et al., 2018).

The presented hypothetical molecular models of BC fate decisions are based on experiments that disentangle the role of various signaling molecules. The simulation results are compared to a mixture of known qualitative and quantitative GC properties. The parameters of the model, in parts, are correlated in the sense that one parameter can compensate for a change of another. For example, Figure S9 (see also the section [Interdependence of parameters](#) in the [STAR Methods](#)) shows how the suppression of pMHC-dependent interaction times between BCs and Tfh cells can be compensated for by parameters controlling DND—at the price of losing physiological relevance, however. In [Figures S11–S16](#), the sensitivity of the parameters in [Table 1](#) is visualized. It can be seen that on the one hand, all of parameters affect the presented readouts, and on the other hand, DND-related parameters have the strongest impact. Thus, while there is theoretical evidence from analysis of experimental data for a DND per selection round on the order of 2 in normal GC reactions (Meyer-Hermann et al., 2012), this was not experimentally proven and a different value likely affects the presented results. The fact that an alternative or even better parameter set may exist for

each model precludes a quantitative weighing of the best consistency of any of the models with real GCs—thus, a quantitative interpretation of the results. The value of this mathematical investigation (Ganusov, 2016) lies in narrowing down the possible roles of molecules by the available experiments and emphasizing the failure of models that do not separate signals for fate decision and division to explain all of the available data. The degree of detail of the proposed molecular pathways was limited to a minimum to limit the size of the parameter space. This implies that pathways are representative of a particular function in the GC reaction and may differ in reality or may be modulated by molecules not considered in this investigation.

STAR★METHODS

Detailed methods are provided in the online version of this paper and include the following:

- [KEY RESOURCES TABLE](#)
- [RESOURCE AVAILABILITY](#)
  - Lead contact
  - Materials availability
  - Data and code availability
- [EXPERIMENTAL MODEL AND SUBJECT DETAILS](#)
- [METHOD DETAILS](#)
  - Germinal center simulation model
  - Space representation
  - Shape space for antibodies
  - B cell phenotypes
  - Founder cells
  - Antigen presentation by FDCs
  - DZ B cell division
  - LZ B cell selection
  - LZ BCs can be in the states unselected, FDC-contact, FDC-selected, Tfh-contact, selected, apoptotic
  - Chemokine distribution
  - Chemotaxis
  - Some general constraints to respect
  - Interdependence of parameters
- [QUANTIFICATION AND STATISTICAL ANALYSIS](#)

SUPPLEMENTAL INFORMATION

Supplemental information can be found online at <https://doi.org/10.1016/j.celrep.2021.109552>.

ACKNOWLEDGMENTS

M.M.H. thanks Mark J. Shlomchik and Gabriel D. Victora for fruitful discussions; Sahamoddin Khailaie for preparation of the graphical abstract and Figure 1 and revision of the manuscript; Sebastian C. Binder, Coraline Mlynarczyk, and Ari M. Melnick for revision of the manuscript; and O. Bannard for suggesting the monitoring of LZ passage times for low- and high-affinity BCs. M.M.H. was in part supported by the Human Frontier Science Program (RGP0033/2015).

AUTHOR CONTRIBUTIONS

M.M.H. developed the research idea and the approach, programmed the code, interpreted the results, and wrote the manuscript.

DECLARATION OF INTERESTS

The author declares no competing interests.

Received: September 29, 2020

Revised: April 6, 2021

Accepted: July 27, 2021

Published: August 24, 2021

REFERENCES

- Allen, C.D., Okada, T., Tang, H.L., and Cyster, J.G. (2007). Imaging of germinal center selection events during affinity maturation. *Science* 315, 528–531.
- Arulraj, T., Binder, S.C., and Meyer-Hermann, M. (2021). Rate of immune complex cycling in follicular dendritic cells determines the extent of protecting antigen integrity and availability to germinal center B cells. *J. Immunol.* 206, 1436–1442.
- Bache, S., and Wickham, H. (2020). *magrittr: a forward-pipe operator for r*. <https://cran.r-project.org/web/packages/magrittr/index.html>.
- Berek, C., and Milstein, C. (1987). Mutation drift and repertoire shift in the maturation of the immune response. *Immunol. Rev.* 96, 23–41.
- Binder, S.C., and Meyer-Hermann, M. (2016). Implications of intravital imaging of murine germinal centers on the control of B cell selection and division. *Front. Immunol.* 7, 593.
- Brandtzaeg, P. (1996). The B-cell development in tonsillar lymphoid follicles. *Acta Otolaryngol. Suppl.* 523, 55–59.
- Brunet, A., Bonni, A., Zigmond, M.J., Lin, M.Z., Juo, P., Hu, L.S., Anderson, M.J., Arden, K.C., Blenis, J., and Greenberg, M.E. (1999). Akt promotes cell survival by phosphorylating and inhibiting a Forkhead transcription factor. *Cell* 96, 857–868.
- Burgering, B.M. (2008). A brief introduction to FOXOlogy. *Oncogene* 27, 2258–2262.
- Calado, D.P., Sasaki, Y., Godinho, S.A., Pellerin, A., Köchert, K., Sleckman, B.P., de Alborán, I.M., Janz, M., Rodig, S., and Rajewsky, K. (2012). The cell-cycle regulator c-Myc is essential for the formation and maintenance of germinal centers. *Nat. Immunol.* 13, 1092–1100.
- Camacho, S.A., Kosco-Vilbois, M.H., and Berek, C. (1998). The dynamic structure of the germinal center. *Immunol. Today* 19, 511–514.
- Chaturvedi, A., Martz, R., Dorward, D., Waisberg, M., and Pierce, S.K. (2011). Endocytosed BCRs sequentially regulate MAPK and Akt signaling pathways from intracellular compartments. *Nat. Immunol.* 12, 1119–1126.
- Cirelli, K.M., Carnathan, D.G., Nogal, B., Martin, J.T., Rodriguez, O.L., Upadhyay, A.A., Enemu, C.A., Gebu, E.H., Choe, Y., Viviano, F., et al. (2020). Slow delivery immunization enhances hiv neutralizing antibody and germinal center responses via modulation of immunodominance. *Cell* 180, 206.
- De Boer, R.J., and Perelson, A.S. (2017). How germinal centers evolve broadly neutralizing antibodies: the breadth of the follicular helper T cell response. *J. Virol.* 91, e00983–17.
- Dengler, H.S., Baracho, G.V., Omori, S.A., Bruckner, S., Arden, K.C., Castrillon, D.H., DePinho, R.A., and Rickert, R.C. (2008). Distinct functions for the transcription factor Foxo1 at various stages of B cell differentiation. *Nat. Immunol.* 9, 1388–1398.
- Depoil, D., Zaru, R., Guiraud, M., Chauveau, A., Harriague, J., Bismuth, G., Utzny, C., Müller, S., and Valitutti, S. (2005). Immunological synapses are versatile structures enabling selective T cell polarization. *Immunity* 22, 185–194.
- Dominguez-Sola, D., Vitorica, G.D., Ying, C.Y., Phan, R.T., Saito, M., Nussenzweig, M.C., and Dalla-Favera, R. (2012). The proto-oncogene MYC is required for selection in the germinal center and cyclic reentry. *Nat. Immunol.* 13, 1083–1091.
- Dominguez-Sola, D., Kung, J., Holmes, A.B., Wells, V.A., Mo, T., Basso, K., and Dalla-Favera, R. (2015). The FOXO1 transcription factor instructs the germinal center dark zone program. *Immunity* 43, 1064–1074.
- Duffy, K.R., Wellard, C.J., Markham, J.F., Zhou, J.H., Holmberg, R., Hawkins, E.D., Hasbold, J., Dowling, M.R., and Hodgkin, P.D. (2012). Activation-induced B cell fates are selected by intracellular stochastic competition. *Science* 335, 338–341.
- Dustin, M.L., and Meyer-Hermann, M. (2012). Immunology. Antigen feast or famine. *Science* 335, 408–409.
- Ersching, J., Efeyan, A., Mesin, L., Jacobsen, J.T., Pasqual, G., Grabner, B.C., Dominguez-Sola, D., Sabatini, D.M., and Vitorica, G.D. (2017). Germinal center selection and affinity maturation require dynamic regulation of mTORC1 kinase. *Immunity* 46, 1045–1058.e6.
- Figge, M.T., Garin, A., Gunzer, M., Kosco-Vilbois, M., Toellner, K.M., and Meyer-Hermann, M. (2008). Deriving a germinal center lymphocyte migration model from two-photon data. *J. Exp. Med.* 205, 3019–3029.
- Fink, S., Hartweg, H., Oliveira, T.Y., Kara, E.E., and Nussenzweig, M.C. (2019). Protein amounts of the MYC transcription factor determine germinal center B cell division capacity. *Immunity* 51, 324–336.e5.
- Ganusov, V.V. (2016). Strong inference in mathematical modeling: a method for robust science in the twenty-first century. *Front. Microbiol.* 7, 1131.
- Garg, A.K., Desikan, R., and Dixit, N.M. (2019). Preferential presentation of high-affinity immune complexes in germinal centers can explain how passive immunization improves the humoral response. *Cell Rep.* 29, 3946–3957.e5.
- Gitlin, A.D., Shulman, Z., and Nussenzweig, M.C. (2014). Clonal selection in the germinal center by regulated proliferation and hypermutation. *Nature* 509, 637–640.
- Hannum, L.G., Haberman, A.M., Anderson, S.M., and Shlomchik, M.J. (2000). Germinal center initiation, variable gene region hypermutation, and mutant B cell selection without detectable immune complexes on follicular dendritic cells. *J. Exp. Med.* 192, 931–942.
- Harrell, F.E.J. (2021). *Hmisc: Harrell miscellaneous*. <https://cran.r-project.org/web/packages/Hmisc/index.html>.
- Hauser, A.E., Junt, T., Mempel, T.R., Sneddon, M.W., Kleinstein, S.H., Henrickson, S.E., von Andrian, U.H., Shlomchik, M.J., and Haberman, A.M. (2007). Definition of germinal-center B cell migration in vivo reveals predominant intrazonal circulation patterns. *Immunity* 26, 655–667.
- Heinzel, S., Binh Giang, T., Kan, A., Marchingo, J.M., Lye, B.K., Corcoran, L.M., and Hodgkin, P.D. (2017). A Myc-dependent division timer complements a cell-death timer to regulate T cell and B cell responses. *Nat. Immunol.* 18, 96–103.
- Hollmann, C., and Gerdes, J. (1999). Follicular dendritic cells and T cells: nurses and executioners in the germinal center reaction. *J. Pathol.* 189, 147–149.
- Hollowood, K., and Macartney, J. (1992). Cell kinetics of the germinal center reaction—a stathmokinetic study. *Eur. J. Immunol.* 22, 261–266.
- Hur, D.Y., Kim, D.J., Kim, S., Kim, Y.I., Cho, D., Lee, D.S., Hwang, Y., Bae, K., Chang, K.Y., and Lee, W.J. (2000). Role of follicular dendritic cells in the apoptosis of germinal center B cells. *Immunol. Lett.* 72, 107–111.
- Inoue, T., Shinnakasu, R., Ise, W., Kawai, C., Egawa, T., and Kurosaki, T. (2017). The transcription factor Foxo1 controls germinal center B cell proliferation in response to T cell help. *J. Exp. Med.* 214, 1181–1198.
- ISO (2012). *ISO/IEC 14882:2011 information technology — programming languages — c++*. <https://www.iso.org/standard/50372.html>.
- Kassambara, A. (2020). *ggpubr: 'ggplot2' based publication ready plots*. <https://cran.r-project.org/web/packages/ggpubr/index.html>.
- Kato, Y., Abbott, R.K., Freeman, B.L., Haupt, S., Groschel, B., Silva, M., Menis, S., Irvine, D.J., Schief, W.R., and Crotty, S. (2020). Multifaceted Effects of Antigen Valency on B Cell Response Composition and Differentiation In Vivo. *Immunity* 53, 548–563.e8.
- Katz, D.H., Paul, W.E., Goidl, E.A., and Benacerraf, B. (1970). Carrier function in anti-hapten immune responses. I. Enhancement of primary and secondary anti-hapten antibody responses by carrier preimmunization. *J. Exp. Med.* 132, 261–282.

- Kuraoka, M., Schmidt, A.G., Nojima, T., Feng, F., Watanabe, A., Kitamura, D., Harrison, S.C., Kepler, T.B., and Kelsoe, G. (2016). Complex antigens drive permissive clonal selection in germinal centers. *Immunity* 44, 542–552.
- Lindhout, E., Lakeman, A., and de Groot, C. (1995). Follicular dendritic cells inhibit apoptosis in human B lymphocytes by a rapid and irreversible blockade of preexisting endonuclease. *J. Exp. Med.* 181, 1985–1995.
- Liu, Y.J., Zhang, J., Lane, P.J., Chan, E.Y., and MacLennan, I.C.M. (1991). Sites of specific B cell activation in primary and secondary responses to T cell-dependent and T cell-independent antigens. *Eur. J. Immunol.* 21, 2951–2962.
- Liu, D., Xu, H., Shih, C., Wan, Z., Ma, X., Ma, W., Luo, D., and Qi, H. (2015). T-B cell entanglement and ICOSL-driven feed-forward regulation of germinal center reaction. *Nature* 517, 214–218.
- Luo, W., Weisel, F., and Shlomchik, M.J. (2018). B Cell Receptor and CD40 Signaling Are Rewired for Synergistic Induction of the c-Myc Transcription Factor in Germinal Center B Cells. *Immunity* 48, 313–326.e5.
- MacLennan, I.C.M. (1994). Germinal centers. *Annu. Rev. Immunol.* 12, 117–139.
- Meyer-Hermann, M. (2014). Overcoming the dichotomy of quantity and quality in antibody responses. *J. Immunol.* 193, 5414–5419.
- Meyer-Hermann, M. (2019). Injection of antibodies against immunodominant epitopes tunes germinal centers to generate broadly neutralizing antibodies. *Cell Rep.* 29, 1066–1073.e5.
- Meyer-Hermann, M., Deutsch, A., and Or-Guil, M. (2001). Recycling probability and dynamical properties of germinal center reactions. *J. Theor. Biol.* 210, 265–285.
- Meyer-Hermann, M.E., Maini, P.K., and Iber, D. (2006). An analysis of B cell selection mechanisms in germinal centers. *Math. Med. Biol.* 23, 255–277.
- Meyer-Hermann, M., Mohr, E., Pelletier, N., Zhang, Y., Victora, G.D., and Toellner, K.M. (2012). A theory of germinal center B cell selection, division, and exit. *Cell Rep.* 2, 162–174.
- Meyer-Hermann, M., Binder, S.C., Mesin, L., and Victora, G.D. (2018). Computer simulation of multi-color Brainbow staining and clonal evolution of B cells in germinal centers. *Front. Immunol.* 9, 2020.
- Miller, M.J., Wei, S.H., Parker, I., and Cahalan, M.D. (2002). Two-photon imaging of lymphocyte motility and antigen response in intact lymph node. *Science* 296, 1869–1873.
- Nossal, G.J. (1992). The molecular and cellular basis of affinity maturation in the antibody response. *Cell* 68, 1–2.
- Omori, S.A., Cato, M.H., Anzelon-Mills, A., Puri, K.D., Shapiro-Shelef, M., Calame, K., and Rickert, R.C. (2006). Regulation of class-switch recombination and plasma cell differentiation by phosphatidylinositol 3-kinase signaling. *Immunity* 25, 545–557.
- Oprea, M., and Perelson, A.S. (1997). Somatic mutation leads to efficient affinity maturation when centrocytes recycle back to centroblasts. *J. Immunol.* 158, 5155–5162.
- Oprea, M., van Nimwegen, E., and Perelson, A.S. (2000). Dynamics of one-pass germinal center models: implications for affinity maturation. *Bull. Math. Biol.* 62, 121–153.
- Papa, I., Saliba, D., Ponzoni, M., Bustamante, S., Canete, P.F., Gonzalez-Figueroa, P., McNamara, H.A., Valvo, S., Grimbaldston, M., Sweet, R.A., et al. (2017). T_{FH}-derived dopamine accelerates productive synapses in germinal centers. *Nature* 547, 318–323.
- Perelson, A.S., and Oster, G.F. (1979). Theoretical studies of clonal selection: minimal antibody repertoire size and reliability of self-non-self discrimination. *J. Theor. Biol.* 81, 645–670.
- R Development Core Team (2020). R: A language and environment for statistical computing (R Foundation for Statistical Computing).
- Rajewsky, K., Schirmacher, V., Nase, S., and Jerne, N.K. (1969). The requirement of more than one antigenic determinant for immunogenicity. *J. Exp. Med.* 129, 1131–1143.
- Robert, P.A., Marschall, A.L., and Meyer-Hermann, M. (2018). Induction of broadly neutralizing antibodies in Germinal Centre simulations. *Curr. Opin. Biotechnol.* 51, 137–145.
- Sander, S., Chu, V.T., Yasuda, T., Franklin, A., Graf, R., Calado, D.P., Li, S., Imami, K., Selbach, M., Di Virgilio, M., et al. (2015). PI3 Kinase and FOXO1 Transcription Factor Activity Differentially Control B Cells in the Germinal Center Light and Dark Zones. *Immunity* 43, 1075–1086.
- Schwicker, T.A., Lindquist, R.L., Shakhar, G., Livshits, G., Skokos, D., Kosco-Vilbois, M.H., Dustin, M.L., and Nussenzweig, M.C. (2007). In vivo imaging of germinal centres reveals a dynamic open structure. *Nature* 446, 83–87.
- Silver, J., Zuo, T., Chaudhary, N., Kumari, R., Tong, P., Giguere, S., Granato, A., Donthula, R., Devereaux, C., and Wesemann, D.R. (2018). Stochasticity enables BCR-independent germinal center initiation and antibody affinity maturation. *J. Exp. Med.* 215, 77–90.
- Stoler-Barak, L., Biram, A., Davidzohn, N., Addadi, Y., Golani, O., and Shulman, Z. (2019). B cell dissemination patterns during the germinal center reaction revealed by whole-organ imaging. *J. Exp. Med.* 216, 2515–2530.
- Tam, H.H., Melo, M.B., Kang, M., Pelet, J.M., Ruda, V.M., Foley, M.H., Hu, J.K., Kumari, S., Crampton, J., Baldeon, A.D., et al. (2016). Sustained antigen availability during germinal center initiation enhances antibody responses to vaccination. *Proc. Natl. Acad. Sci. USA* 113, E6639–E6648.
- Tas, J.M., Mesin, L., Pasqual, G., Targ, S., Jacobsen, J.T., Mano, Y.M., Chen, C.S., Weill, J.C., Reynaud, C.A., Browne, E.P., et al. (2016). Visualizing antibody affinity maturation in germinal centers. *Science* 351, 1048–1054.
- Tew, J.G., Wu, J., Qin, D., Helm, S., Burton, G.F., and Szakal, A.K. (1997). Follicular dendritic cells and presentation of antigen and costimulatory signals to B cells. *Immunol. Rev.* 156, 39–52.
- Thaunat, O., Granja, A.G., Barral, P., Filby, A., Montaner, B., Collinson, L., Martinez-Martin, N., Harwood, N.E., Bruckbauer, A., and Batista, F.D. (2012). Asymmetric segregation of polarized antigen on B cell division shapes presentation capacity. *Science* 335, 475–479.
- Toellner, K.M., Jenkinson, W.E., Taylor, D.R., Khan, M., Sze, D.M.Y., Sansom, D.M., Vinuesa, C.G., and MacLennan, I.C.M. (2002). Low-level hypermutation in T cell-independent germinal centers compared with high mutation rates associated with T cell-dependent germinal centers. *J. Exp. Med.* 195, 383–389.
- Turner, J.S., Ke, F., and Grigorova, I.L. (2018). B cell receptor crosslinking augments germinal center B cell selection when t cell help is limiting. *Cell Rep.* 25, 1395–1403.e4.
- Tzivion, G., Dobson, M., and Ramakrishnan, G. (2011). FoxO transcription factors: regulation by AKT and 14-3-3 proteins. *Biochim. Biophys. Acta* 1813, 1938–1945.
- van Eijk, M., Defrance, T., Hennino, A., and de Groot, C. (2001). Death-receptor contribution to the germinal-center reaction. *Trends Immunol.* 22, 677–682.
- Victora, G.D., and Nussenzweig, M.C. (2012). Germinal centers. *Annu. Rev. Immunol.* 30, 429–457.
- Victora, G.D., Schwicker, T.A., Fooksman, D.R., Kamphorst, A.O., Meyer-Hermann, M., Dustin, M.L., and Nussenzweig, M.C. (2010). Germinal center dynamics revealed by multiphoton microscopy with a photoactivatable fluorescent reporter. *Cell* 143, 592–605.
- Victora, G.D., Dominguez-Sola, D., Holmes, A.B., Deroubaix, S., Dalla-Favera, R., and Nussenzweig, M.C. (2012). Identification of human germinal center light and dark zone cells and their relationship to human B-cell lymphomas. *Blood* 120, 2240–2248.
- Wan, Z., Lin, Y., Zhao, Y., and Qi, H. (2019). T_{FH} cells in bystander and cognate interactions with B cells. *Immunol. Rev.* 288, 28–36.
- Wang, S., Mata-Fink, J., Kriegsmann, B., Hanson, M., Irvine, D.J., Eisen, H.N., Burton, D.R., Wittrup, K.D., Kardar, M., and Chakraborty, A.K. (2015). Manipulating the selection forces during affinity maturation to generate cross-reactive HIV antibodies. *Cell* 160, 785–797.
- Wang, P., Shih, C.M., Qi, H., and Lan, Y.H. (2016). A stochastic model of the germinal center integrating local antigen competition, individualistic T-B interactions, and B cell receptor signaling. *J. Immunol.* 197, 1169–1182.

- Wickham, H. (2007). Reshaping data with the reshape package. *J. Stat. Softw.* 21, 1–20.
- Wickham, H. (2016). *ggplot2: Elegant Graphics for Data Analysis* (Springer).
- Wilhelm, K., Happel, K., Eelen, G., Schoors, S., Oellerich, M.F., Lim, R., Zimmermann, B., Aspalter, I.M., Franco, C.A., Boettger, T., et al. (2016). FOXO1 couples metabolic activity and growth state in the vascular endothelium. *Nature* 529, 216–220.
- Zhang, J., and Shakhnovich, E.I. (2010). Optimality of mutation and selection in germinal centers. *PLoS Comput. Biol.* 6, e1000800.
- Zhang, Y., Meyer-Hermann, M., George, L.A., Figge, M.T., Khan, M., Goodall, M., Young, S.P., Reynolds, A., Falciani, F., Waisman, A., et al. (2013). Germinal center B cells govern their own fate via antibody feedback. *J. Exp. Med.* 210, 457–464.
- Zinkernagel, R.M. (1997). Felix Hoppe-Seyler Lecture 1997. Protective antibody responses against viruses. *Biol. Chem.* 378, 725–729.

STAR★METHODS

KEY RESOURCES TABLE

RESOURCE	SOURCE	IDENTIFIER
Software, algorithms, and experimental data		
Simulation software	Zenodo	Database: https://doi.org/10.5281/zenodo.4662037
Data in Figures 3A–3C, 4A	(Victora et al., 2010)	N/A
Data in Figures 5A–5C	(Tas et al., 2016)	N/A
C++ ISO/IEC 14882:2011	(ISO, 2012)	http://www.iso.org/iso/iso_catalogue/catalogue_tc/catalogue_detail.htm?csnumber=50372
R	(R Development Core Team, 2020)	https://www.R-project.org/
R package ggplot2	(Wickham, 2016)	https://ggplot2.tidyverse.org
R package ggpubr	(Kassambara, 2020)	https://cran.r-project.org/web/packages/ggpubr/index.html
R package Hmisc	(Harrell, 2021)	https://cran.r-project.org/web/packages/Hmisc/index.html
R package magrittr	(Bache and Wickham, 2020)	https://cran.r-project.org/web/packages/magrittr/index.html
R package reshape2	(Wickham, 2007)	https://cran.r-project.org/web/packages/reshape2/index.html

RESOURCE AVAILABILITY

Lead contact

Further information and requests can be addressed to Michael Meyer-Hermann (mmh@theoretical-biology.de).

Materials availability

This study did not generate new unique reagents.

Data and code availability

- This paper analyzes existing, publicly available data. References to the datasets are found in the respective figure legends and are listed again in the [Key resources table](#).
- All original code has been deposited at Zenodo and is publicly available as of the date of publication. DOIs are listed in the [Key resources table](#).
- Post-publication availability of data and code

Any additional information required to reanalyze the data reported in this paper is available from the lead contact upon request.

EXPERIMENTAL MODEL AND SUBJECT DETAILS

This study did not involve any experiments and exclusively relies on computer simulations.

METHOD DETAILS

Germinal center simulation model

The germinal center (GC) model assumptions underlying the GC simulations are explained here and include the used parameter values. It follows [Meyer-Hermann et al. \(2012\)](#) with additional features ([Meyer-Hermann, 2014](#); [Binder and Meyer-Hermann, 2016](#); [Meyer-Hermann, 2019](#)), and adding the specificities of the DisseD-, the BCinTime-, and the MiXed-models introduced here. Parameters differing in the three theories are listed in [Table 1](#). Used acronyms are: DZ for dark zone, LZ for light zone, Tfh for T follicular helper, FDC for follicular dendritic cell, BC for B cell, BCR for B cell receptor.

Space representation

All reactions take place on a three-dimensional discretized space with a rectangular lattice with lattice constant of $\Delta x = 5\mu m$. Every lattice node can be occupied by a single cell only.

Shape space for antibodies

Antibodies are represented on a four ($d = 4$) dimensional shape space (Perelson and Oster, 1979). The shape space is restricted to a size of 10 positions per dimension, thus, only considering antibodies with a minimum affinity to the antigen. The optimal clone Φ^* is positioned in the center of the shape space. A position on the shape space Φ is attributed to each BC.

The 1-Norm with respect to the optimal clone $\|\Phi - \Phi^*\|_1 = \sum_{i=1}^d |\Phi_i - \Phi_i^*|$, i.e., the minimum number of mutations required to reach the optimal clone, is used as a measure for the antigen binding probability. The binding probability is calculated from the Gaussian distribution with width $\Gamma = 2.8$ (Meyer-Hermann et al., 2001):

$$b(\Phi, \Phi^*) = \exp\left(-\frac{\|\Phi - \Phi^*\|_1^2}{\Gamma^2}\right). \quad (\text{Equation 1})$$

B cell phenotypes

Three BC phenotypes are distinguished: DZ BCs, LZ BCs, and output cells. The different phenotypes characterize the cell properties and are not meant as localization within the GC zones. DZ BCs divide, mutate and migrate. LZ BCs also migrate and undergo the different stages of the selection process. Output cells only migrate.

Founder cells

The model starts from a fixed number of Tfh cells (Table 1), 200 FDCs, 300 stromal cells, and no BC. Tfh cells are randomly distributed on the lattice and occupy a single node each. Stromal cells are restricted to the DZ (see section Chemokine distribution for their function). FDCs are restricted to the upper half of the reaction sphere, occupy one node by their soma and have 6 dendrites of $40\mu m$ length each. The presence of dendrites is represented as a lattice-node property and, thus, visible to BCs. The dendrites are treated as transparent for BC or Tfh cell migration such that they do not inhibit cell motility.

BCs enter the GC reaction with a probability per time step corresponding to a rate of 2 cells per hour (for an estimation of this value refer to Meyer-Hermann, 2019). New BCs are randomly positioned on free lattice nodes. The shape space position of each new BC is randomly picked from a set of 100 randomly picked shape space positions, unless stated otherwise.

Antigen presentation by FDCs

Each FDC is loaded with antigen portions distributed onto the lattice-nodes occupied by FDC-soma or FDC-dendrite (Table 1). One antigen portion corresponds to the number of antigen molecules taken up by a BC upon successful contact with an FDC.

DZ B cell division

The average cell cycle duration of 7.5 hours of DZ BCs is varied for each BC according to a Gaussian distribution. This is needed to get desynchronization of BC division. The cell cycle is decomposed into four phases (G1, S, G2, M) in order to localize mitotic events if this is needed.

Each founder BC divides a number of times before differentiating to the LZ phenotype for the first time. Six divisions was the number of divisions found in response to the extreme stimulus with anti-DEC205-OVA (Victora et al., 2010; Meyer-Hermann et al., 2012). Each selected BC divides a number of times determined by the interaction with Tfh cells (see below, LZ BC selection). The parameters of the interaction with Tfh cells are tuned such that the mean number of divisions is in the range of two (Gitlin et al., 2014). This value is required in order to maintain a DZ to LZ ratio in the range of two (Victora et al., 2010; Meyer-Hermann et al., 2012).

A division requires free space on one of the Moore neighbors of the dividing cell. Otherwise the division is postponed until a free Moore neighbor is available.

At every division the encoded antibody mutates with probability 0.5 (Berek and Milstein, 1987; Nossal, 1992). This corresponds to a shift in the shape space to a von Neumann neighbor in a random direction. Upon selection by Tfh cells the mutation probability is individually reduced from $m_{\max} = 0.5$ down to $m_{\min} = 0$ in an affinity-dependent way following

$$m(b) = m_{\max} - (m_{\max} - m_{\min})b = \frac{1 - b}{2} \quad (\text{Equation 2})$$

with b from Equation 1 (Toellner et al., 2002). Thus, after recycling, DZ BCs can acquire reduced mutation probabilities. This mechanism is motivated by the observation that BCR internalization enhances the activation of the kinase Akt (Chaturvedi et al., 2011) which, in turn, suppresses activation induced cytosine deaminase (AID) (Omori et al., 2006). AID is required for somatic hypermutation, such that this provides an affinity-dependent downregulation of the mutation frequency (Dustin and Meyer-Hermann, 2012).

BC division of BCs that previously acquired antigen and have been selected by Tfh cells distribute the retained antigen asymmetrically to the daughters (Thaunat et al., 2012). The model assumes asymmetric division in 72% of the cases, which is supported by

experimental observations (see [Thaunat et al., 2012](#) and Figure S1 in [Meyer-Hermann et al., 2012](#)). If division is asymmetric, one daughter gets all the retained antigen while the other gets none, which approximates the value of 88% found in ([Thaunat et al., 2012](#)). Mutation is suppressed in cells retaining antigen.

After the required number of divisions, the BC differentiates with a rate of in 1/6 minutes to the LZ phenotype. All BCs that kept the antigen up to this time, differentiate to output cells, upregulate CXCR4, and leave the GC in direction of the T zone.

LZ B cell selection

At the time $t = 0$ of differentiation from the DZ to the LZ phenotype, BCs are in state *unselected* and BC-specific factors are initialized to

$$\begin{aligned} F(t=0) &= F_0 = 1 \\ C(t=0) &= C_0 = 0 \\ R(t=0) &= R_0 = 0, \end{aligned} \quad (\text{Equation 3})$$

where F_0 does not apply to BCinTime.

These factors are associated with FoxO1 (F), c-Myc (C), and mTOR (R) in the DisseD- and the MiXed-models. In BCs not in contact to Tfh cells, FoxO1, mTOR and c-Myc evolve according to

$$\frac{dF}{dt} = -\alpha_F H_F(B) F \quad (\text{Equation 4})$$

$$\frac{dR}{dt} = \beta_R H_R(B) \quad (\text{Equation 5})$$

$$\frac{dC}{dt} = -\gamma_C C \quad (\text{Equation 6})$$

with $\beta_R = 1/(8 \text{ hours})$ the mTOR production rate modulated by the BCR-dependent Hill-function $H_R(B)$ (see K_R and n_R in [Table 1](#) and [Figure S10B](#)). $\alpha_F = 2/\text{hours}$ is the FoxO1 reduction rate modulated by the BCR-dependent Hill-function $H_F(B)$ between one and zero with K-value of 20 and Hill-coefficient 1 ([Figure S10C](#)). $\gamma_C = \ln(2)/(3 \text{ hours})$ is the c-Myc degradation rate. R and C have different dynamics in state *Tfh-contact*.

In BCinTime, the unspecified differentiation signal (R) follows

$$\frac{dR}{dt} = \beta_{\min} + (\beta_{\max} - \beta_{\min}) \frac{p^{n_R}}{p^{n_R} + K_R^{n_R}}, \quad (\text{Equation 7})$$

with $1/\beta_{\min} = 2 \text{ hours}$ and $1/\beta_{\max} = 18 \text{ hours}$. Thus, the growth rate of R is determined by the total amount of pMHC p collected by the BC at any given time with K_R and n_R the pMHC amount for half max reduction of the growth rate and the associated Hill-coefficient, respectively, (see [Table 1](#) and [Figure S10A](#)). The integrated level of Tfh cell signals a BC received (C) is defined in the state *Tfh-contact*, and F is not used.

LZ BCs can be in the states *unselected*, *FDC-contact*, *FDC-selected*, *Tfh-contact*, *selected*, *apoptotic*

Unselected. LZ BCs migrate and search for contact with FDCs loaded with antigen in order to collect antigen for 0.7 hours. If an FDC soma or dendrite is present at the position of the BC, the BC attempts to establish contact to the epitope with highest affinity to the BCR (default setting). Alternatively, the BC may attempt to establish contact to the epitope of highest availability at this site. In both settings, binding is affinity dependent and happens with the probability b in [Equation 1](#). If the available number of antigen portions at the specific FDC site drops below 20 the binding probability b is linearly reduced with the number of available portions. If successful, the BC switches to the state *FDC-contact*; otherwise the BC continues to migrate. Further binding-attempts are prohibited for 1.2 minutes. At the end of the antigen collection period, BCs switch to the state *FDC-selected*. If a LZ BC fails to collect any antigen until this time it switches to the state *apoptotic*.

FDC-contact. LZ BCs remain immobile (bound) for 3 minutes ([Schwickert et al., 2007](#)) and then return to the state *unselected* or *FDC-selected*, depending on from which state contact to FDC was established. The counter for the number of successful antigen uptake events p and the degree of BCR-signaling B are increased by one unit and the FDC reduces its locally available antigen portions by one. In DisseD and MiXed, factors depend differently on the amount of presented pMHC p and the degree of BCR-signaling B . In anti-DEC205-OVA experiments, antigen is taken up and presented without induction of BCR signals, thus, p is increased by 100 antigen uptake events but B is left unchanged. The amount of 100 is motivated by the finding that pMHC presentation is increased five-fold upon anti-DEC205-OVA injection ([Victoria et al., 2010](#)) together with a mean number of 20 – 30 uptake events seen under normal conditions in the simulations.

FDC-selected. BCs search for contact with Tfh cell or FDC. If they meet a Tfh cell, they switch to the state *Tfh-contact*. If they meet a FDC they switch to the state *FDC-contact* with probability b in Equation 1. If Tfh cells and FDCs are neighbors of a BC, the BC binds the Tfh cell.

Tfh-contact. LZ BCs remain immobile for a time randomly sampled from a Gaussian distribution with mean 6 minutes and width 1.2 minutes (BCinTime) or for a time determined by a Hill-function in dependence on the amount of presented pMHC p (DisseD and MiXed), see Table 1 *Tfh-BC-interaction time* and Figure S10F. During this time, the bound Tfh cells, which may also be bound to other BCs, polarizes to the BC with highest number of successful antigen uptakes. Polarization is determined from scratch in every time step and inertia from repolarization or intracellular organelles are ignored. If more than one BC shares the same highest pMHC presentation p , polarization is chosen randomly among those BCs. Only the BC to which the Tfh cell is polarized receives Tfh cell signals C and accumulate those.

In BCinTime, Tfh cell-signals follow

$$\frac{dC}{dt} = \alpha_C \delta_{TB}, \quad (\text{Equation 8})$$

where α_C is the rate of signal acquisition during polarized Tfh cell-BC interactions, ensured by δ_{TB} , which is one during polarized contact and zero otherwise. The rate is set to 1/hr without loss of generality. Thus, C reflects the integrated duration of Tfh cell-BC interactions with Tfh cells polarized to this BC.

In DisseD and MiXed, Tfh cell signals are represented as mTOR levels as well as c-Myc levels, which are modulated by FoxO1, according to

$$\frac{dC}{dt} = \alpha_C (1 - H_C(F)) T(p) H_I(t) \delta_{TB} - \gamma_C C \quad (\text{Equation 9})$$

$$\frac{dR}{dt} = \alpha_R H_R(B) H_I(t) T(p) \delta_{TB} + \beta_R H_R(B). \quad (\text{Equation 10})$$

$T(p)$ is the pMHC-dependent intensity of Tfh cell signaling to the interacting BC associated with CD40-signals, which is described by a Hill-function (see Table 1 $T(p)$ and Figure S10G). Signaling intensity is further modulated by the degree of ICOSL upregulation $H_I(t)$. $H_I(t)$ is a Hill-function in time with values between zero and one, determined by a characteristic ICOSL-upregulation time of four hours in mice (Papa et al., 2017) and Hill-coefficient two (Figure S10E). α_C and γ_C are the growth and degradation rates of c-Myc. $\beta_R = 1/(8 \text{ hours})$ tunes the level of mTOR upregulation when BCs are not in contact to Tfh cells (see Equation 5), while $\alpha_R = 1/(\text{hours})$ determines mTOR upregulation during productive contact with a Tfh cell. $H_R(B)$ is a Hill-function describing the dependence of mTOR regulation on signals downstream of BCR (see K_R and n_R in Table 1 and Figure S10B). $H_C(F)$ is a Hill-function controlling the suppression of c-Myc signals by FoxO1 with values between zero and one, K-value of 0.5 and Hill-coefficient of 1 (Figure S10D).

After the binding time, the BC detaches and returns to the state *FDC-selected*. It continues to search for and bind FDCs or Tfh cells, where binding twice the same Tfh cell in a sequence is excluded. The total period of signal acquisition in the LZ is determined differently in the three theories: It finished when $R=1$ is reached in BCinTime, when $R=1$ or the maximum time period of 18 hours is reached in DisseD, when the LZ passage time gets longer than the number of antigen uptake events times 0.5 hours in MiXed. When one of those limits is reached during a running contact, the contact is kept active until the end of the contact time. Then, it switches to the state *apoptotic* if either $R < 1$ or if the received Tfh cell-signals are $C < C_{\text{threshold}}$. It switches to the state *selected* otherwise.

Selected. LZ BCs keep the LZ phenotype for a determined time period (Table 1, LZ-DZ BC-differentiation delay) and desensitize for CXCL13, thus, perform a random walk. During that time they re-enter cell cycle and progress through the cell cycle phases. Then they recycle back to the DZ phenotype with a rate of 1/6 minutes and memorize the amount of collected antigen as well as the cell cycle phase they have reached by this time.

The number of divisions $P(C)$ the recycled BCs will do is derived from C , which reflects the amount of integrated Tfh cell signals (BCinTime) or the c-Myc level (DisseD and MiXed), according to

$$P(C) = P_{\min} + (P_{\max} - P_{\min}) \frac{C^{n_p}}{C^{n_p} + K_p^{n_p}}. \quad (\text{Equation 11})$$

The minimum number of division is set to one ($P_{\min} = 1$) in order to avoid recycling events without further division. It is limited by six divisions in the best case, which is motivated by anti-DEC205-OVA experiments in which DEC205^{+/+} BCs received abundant antigen which increased pMHC presentation to a maximum (Victora et al., 2010). The population dynamics *in vivo* and *in silico* only matched when the number of divisions was increased to six in the simulation (Meyer-Hermann et al., 2012) suggesting that the strongest possible pMHC presentation to Tfh cells induces six divisions ($P_{\max} = 6$). The Hill-coefficient n_p was model-specific (Table 1 and Figure S10H).

The half value K_P remained to be determined, which denotes the amount of antigen collected by BCs at which the number of divisions becomes half maximal. It was fixed by the condition that for the mean level of $C = C_0$ the number of divisions becomes two, as observed in Gitlin et al. (2014):

$$K_P \approx C_0 \left(\frac{P_{\max} - P_{\min}}{P_0 - P_{\min}} - 1 \right)^{1/n_P} \quad (\text{Equation 12})$$

and differs for each model (Table 1).

Apoptotic. LZ BCs remain on the lattice for 6 hours before they are deleted. They continue to be sensitive to CXCL13 during this time.

Chemokine distribution

Two chemokines CXCL12 and CXCL13 are considered. CXCL13 is produced by FDCs in the LZ with 10nMol per hour and FDC while CXCL12 is produced by stromal cells in the DZ with 400nMol per hour and stromal cell. As both cell types are assumed to be immobile, chemokine distributions were pre-calculated once and the resulting steady state distributions were used in all simulations.

Chemotaxis

DZ and LZ BCs regulate their sensitivity to CXCL13 and CXCL12, respectively. This is true in all BC states unless stated otherwise. All BCs move with a target speed of $7.5 \mu\text{m}/\text{min}$. This leads to a slightly lower observable average speed of $\approx 6 \mu\text{m}/\text{min}$.

BCs have a polarity vector that determines their preferential direction of migration. The polarity vector $\vec{p}$ is reset every 1.5 minutes into a new direction using the chemokine distribution c as

$$\vec{p} = \vec{p}_{\text{rand}} + \frac{\alpha}{1 + \exp\left\{ \kappa \left(K_{1/2} - \Delta x |\vec{\nabla} c| \right) \right\}} \frac{\vec{\nabla} c}{|\vec{\nabla} c|}, \quad (\text{Equation 13})$$

where $\vec{p}_{\text{rand}}$ is a random polarity vector and the turning angle is sampled from the measured turning angle distribution (Allen et al., 2007 Figure S1B). $\alpha = 3$ determines the relative weight of the chemotaxis and random walk, $K_{1/2} = 2 \cdot 10^{11}$ Mol determines the gradient of half maximum chemotaxis weight, and $\kappa = 10^{10}$ /Mol determines the steepness of the weight increase.

BCs de- and re-sensitize for their respective chemokine depending on the local chemokine concentration: The desensitization threshold is set to 4.5nMol and 0.08nMol for CXCL12 and CXCL13, respectively, which avoids cell clustering in the center of the zones. The resensitization threshold is set at 2/3 and 3/4 of the desensitization threshold for CXCL12 and CXCL13, respectively.

BCs can only migrate if the target node is free. If occupied and the neighbor cell is to migrate in the opposite direction (negative scalar product of the polarity vectors) both cells are exchanged with a probability of 0.5. This exchange algorithm avoids lattice artifacts leading to cell clusters.

Tfh cells do random walk with a preferential directionality to the LZ: The polarity vector $\vec{p}$ of Tfh cells is determined from a mixture of random walk $\vec{r}$ and the direction of the LZ $\vec{n}$ by

$$\vec{p} = (1 - \alpha') \vec{r} + \alpha' \vec{n}, \quad (\text{Equation 14})$$

where $\alpha' = 0.1$ is the weight of chemotaxis. This weight leads to a dominance of random walk with a tendency to accumulate in the LZ as found in experiment. Tfh cells migrate with an average speed of $10 \mu\text{m}/\text{min}$ and repolarize every 1.7 minutes (Miller et al., 2002).

Output cell motility is derived from plasma cell motility data to $3 \mu\text{m}$ per minute with a persistence time of 0.75 minutes (Allen et al., 2007).

Some general constraints to respect

There is a number of qualitative features that any GC simulation needs to respect. Some important such features are listed here.

- GC kinetics must exhibit a rising phase, a peak between day 6 and 9 post GC onset, and then a declining phase. The overall probability of death for high affinity BCs must be higher than the probability of selection and sets boundaries on parameters associated with DND and apoptosis. On the long run, antigen consumption further reduces the probability of selection and DND.
- Affinity maturation needs to lead to a relevant amount of high affinity cells at day 8 post GC onset. This enforces a comparably high selection pressure in all models.
- In contrast to BCinTime, in DisseD, high affinity BC are selected faster than low affinity BCs. Therefore, mTOR must be upregulated faster in high affinity cells as guaranteed by the $H_R(B)T(p)$ term in Equation 10, which depends on BCR-signals B and pMHC-presentation p .
- Selected high affinity BCs do more divisions in the DZ (Gitlin et al., 2014), i.e., high affinity cells must achieve a higher c-Myc score. This is reflected in the $T(p)$ term in Equation 9. However, a higher c-Myc score has still to be achieved even though the low affinity cells stay longer in the LZ, thus, having more time for BC-Tfh cell-interactions. Competition for Tfh cell help in a

pMHC-dependent way might not be sufficient. This speaks for a decay of c-Myc on a timescale comparable to the time between two BC-Tfh cell-interactions, which would require an efficient and ongoing uptake of antigen via the BCR in order to achieve a high c-Myc level at the time of selection. This is realized with the $\gamma_C C$ term in Equation 9.

- The achieved level of C at the time of selection $R = 1$ should induce a DND of less than 2 for low and between 2 and 4 for high affinity BCs. The total mean of DND should remain in the range of 2 (Meyer-Hermann et al., 2012). This determines the steepness of the DND function Equation 11 and the half value K_P .
- c-Myc upregulation should require both, BCR signaling and Tfh cell help (Luo et al., 2018). This is realized by the introduction of FoxO1-dynamics (Equation 4) and the product $(1 - H_C(F))T(p)$ in Equation 9. The FoxO1-brake is released dependent on BCR signals and c-Myc upregulation requires the released FoxO1-brake plus Tfh cell signals in a pMHC-dependent manner.
- It should be possible that antigen-uptake by BCR alone can lead to selection, even without Tfh cell help. This implies that mTOR is able to grow even without Tfh cell signals in Equation 10 and requires the additive term $\beta_R H_R(B)$, such that even with $T = 0$ a remaining purely BCR signaling-dependent mTOR signal is possible. However, β_R should remain small in comparison to the term dependent on $T(p)$ for intermediate pMHC levels.
- It should be possible to get selected with Tfh cell help alone with minor BCR signals. This would recapitulate the anti-DEC205-OVA experiments (Victora et al., 2010) and imply that a small BCR signal is sufficient for upregulation of mTOR. As BCs that received antigen via their DEC205-receptor would continue collecting antigen, even though in a rather inefficient way for most of the low affinity cells, a small BCR signal might be there allowing for the product $H_R(B)T(p)$ in Equation 10 to generate some mTOR signal.
- High pMHC but low BCR should lead to a prolongation of the LZ phase until selection (Victora et al., 2010; Meyer-Hermann et al., 2012). This implies that mTOR upregulation is slow despite high pMHC-dependent Tfh cell signals and is reflected in the low $H_R(B)$ value in Equation 10, which strongly delays reaching $R = 1$. The relative K-values in the Hill-functions $H_{F,R}$ in Equations 4 and 10 should respect $K_F < K_R$ in order to allow for a release of the FoxO1-brake on c-Myc at low BCR signals while keeping a slow upregulation of mTOR.
- High pMHC but low BCR should lead to high c-Myc scores reflecting the strong DND observed in the anti-DEC205-OVA experiment (Victora et al., 2010). Thus, c-Myc should go up despite an ongoing suppression by FoxO1, which is only released by BCR signals. This implies that in the low BCR signaling limit, suppression of c-Myc by FoxO1 is weaker than suppression of mTOR by lack of BCR signals, again determining the relative values of $K_F < K_R$ in Equations 4 and 10.

Interdependence of parameters

The present investigation validates mathematical models with a mixture of quantitative and qualitative experimental data. This does not allow for the definition of a quantitative cost function to systematically evaluate the quality of the respective model. In particular, correlations between pairs of parameters are likely to exist.

In order to still illustrate the interdependence of parameters, the particular example of pMHC-dependent duration of BC-Tfh cell-interactions was considered, which is active in DisseD and MiXed. In both models, the duration of every BC-Tfh cell-interaction was set constant. This implies that help signals from Tfh cells to BCs remain the same, irrespective of the density of pMHC presented on the BCs (which reflects its affinity for the antigen). The resulting GCs were substantially smaller (see Figure S9): In the beginning, selection pressure was weak because even low affinity BCs received signals from Tfh cells for the now fixed period of time. This delayed affinity maturation and induced an intermediate shrinking of the GC population because high affinity cells did not receive sufficient advantage. When sufficient high affinity BCs were generated, an unphysiological second wave in the GC population evolved.

The lower GC size was a result of reduced mean DND (see Figure S9). This could be partly compensated by a reduction of the DND-parameter K_P , which makes DND more sensitive to the lower overall c-Myc level. However, affinity maturation remained delayed, which is not supported by experiment. Further, the burst after application of anti-DEC205-OVA did not match experimental results to the same extent (see Figure S9, right panels).

This example illustrates that Tfh cell-signal strength and DND parameters are correlated, but that this involves more than one parameter. The author did not find a parameter set that matched experimental constraints to the same degree as the models in the main text. This supports that the dynamics of Tfh cell-signals are important to keep consistency with experiment. Note that there are many partially inter-related parameters.

QUANTIFICATION AND STATISTICAL ANALYSIS

All graphs were generated with R. The corresponding R files are part of the data at Zenodo (see Key resources table). All simulations were repeated 100 or 1000 times (see Figure legends) with random number generator seeded by the CPU time. Mean and standard deviation are shown in the figures.