

Tables

Table 1. Incidence of anogenital warts in the age group 10-79 years in Germany from 2005 - 2008

	Year	Person-time (in 100,000 years)	Cases	Incidence (per 100,000 person years)	95% Confidence Interval
Total	2005	45.70	7662	167.66	[163.92; 171.67]
	2006	47.99	8190	170.67	[166.99; 174.40]
	2007	48.17	8729	181.21	[177.43; 185.06]
	2008	51.78	10,169	196.38	[192.59; 200.24]
Male	2005	24.27	3270	134.76	[130.18; 139.46]
	2006	25.28	3561	140.87	[136.28; 145.58]
	2007	25.37	3874	152.68	[147.91; 157.57]
	2008	26.84	4590	171.02	[166.11; 176.04]
Female	2005	21.44	4392	204.90	[198.88; 211.05]
	2006	22.71	4629	203.83	[198.00; 209.79]
	2007	22.80	4855	212.97	[207.02; 219.05]
	2008	24.94	5579	223.68	[217.85; 229.63]

Table 2. Changes in the incidence of anogenital warts among females in the age group 12 to 19 years in 2005-2008 (separate model in each line, with 2nd quarter of 2007 as the breaking point^a)

Age [years]	Mean number of cases* per quarter	Range	Mean person time per quarter ^b	Relative change of incidence per quarter ^c	95% Confidence interval	p-value	Estimated reduction in the incidence per year
12	0.9	[0;3]	8.47	0.79	[0.54; 1.14]	0.211	61.36%
13	1.3	[0;3]	8.41	1.13	[0.92; 1.40]	0.252	-63.95%
14	4.4	[1;10]	8.32	1.07	[0.96; 1.19]	0.248	-28.76%
15	8.9	[4;17]	8.05	0.99	[0.91; 1.07]	0.768	4.80%
16	13.9	[5;25]	7.67	0.90	[0.84; 0.97]	0.004	34.79%
17	19.3	[9;28]	7.26	0.91	[0.85; 0.96]	0.001	32.75%
18	28.5	[21;35]	7.06	0.93	[0.89; 0.98]	0.005	24.66%
19	37.4	[27;49]	7.21	0.98	[0.98; 1.02]	0.221	9.48%

* persons with new diagnoses of anogenital warts

^a additionally adjusted for quarters within the year

^b in the cohort in 1,000 person years

^c after the breaking point (risk ratio, adjusted for overall trend and seasonality)