

Draft Genome Sequence of the Deep-Subsurface Actinobacterium *Tessaracoccus lapidicaptus* IPBSL-7^T

Fernando Puente-Sánchez,^a Dietmar H. Pieper,^b Alejandro Arce-Rodríguez^{b*}

Systems Biology Program, Centro Nacional de Biotecnología, Consejo Superior de Investigaciones Científicas (CSIC), Madrid, Spain^a; Microbial Interactions and Processes Research Group, Helmholtz Centre for Infection Research, Braunschweig, Germany^b

* Present address: Alejandro Arce-Rodríguez, Molecular Bacteriology Group, Helmholtz Centre for Infection Research, Braunschweig, Germany.

The type strain of *Tessaracoccus lapidicaptus* was isolated from the deep subsurface of the Iberian Pyrite Belt (southwest Spain). Here, we report its draft genome, consisting of 27 contigs with a ~3.1-Mb genome size. The annotation revealed 2,905 coding DNA sequences, 45 tRNA genes, and three rRNA genes.

Received 9 August 2016 Accepted 10 August 2016 Published 29 September 2016

Citation Puente-Sánchez F, Pieper DH, Arce-Rodríguez A. 2016. Draft genome sequence of the deep-subsurface actinobacterium *Tessaracoccus lapidicaptus* IPBSL-7^T. *Genome Announc* 4(5):e01078-16. doi:10.1128/genomeA.01078-16.

Copyright © 2016 Puente-Sánchez et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Fernando Puente-Sánchez, fpusan@gmail.com.

Members of the genus *Tessaracoccus* are Gram-positive, non-motile, nonsporulating, and facultative anaerobic actinobacteria (1). Two species, *T. profundus* and *T. lapidicaptus*, have been isolated from deep-subsurface samples (2, 3). *T. lapidicaptus* was recently isolated from a 297-m-deep sample obtained from a drilling project that aimed to intersect the groundwater interacting with the massive pyrite minerals present in the Iberian Pyrite Belt (southwest Spain) subsurface (4). This environment is of special interest, as underground microbial oxidation of pyrite has been hypothesized to be responsible for the unique characteristics of Río Tinto (“Red River”), a natural extremely acidic stream with high concentrations of heavy metals (5). Furthermore, *T. lapidicaptus* possesses bioremediation potential due to its ability to mediate the biological precipitation of iron-rich phosphates and carbonates under anaerobic conditions (6). The study of its genome may help to identify genes involved in iron biomineralization and heavy metal resistance and to elucidate the particular adaptations that allow this microorganism to thrive under the extreme energetic and nutritional limitations that are characteristic of deep-subsurface environments (7).

A lyophilized sample of *T. lapidicaptus* IPBSL-7^T (DSMZ-27266) was obtained from the German Collection of Microorganisms and Cell Cultures (DSMZ, Braunschweig, Germany). Briefly, the cell pellet was rehydrated and cultured for 2 days in 5 ml of tryptic soy broth at 30°C. Total genomic DNA was isolated from 2 ml of the culture using the DNeasy blood and tissue kit (Qiagen, Düsseldorf, Germany). After DNA shearing (Covaris, Woburn, MA, USA), Illumina paired-end libraries were prepared using the NEBNext Ultra DNA library prep kit (NEB, Ipswich, MA, USA) and subjected to 250-bp paired-end sequencing on an Illumina MiSeq platform (Illumina, San Diego, CA, USA), which generated a total of 1,989,155 read pairs. Adaptor sequences were removed with Cutadapt version 1.10 (8), and reads were quality-trimmed by using PRINSEQ-lite version 0.20.4 (9). The resulting 1,245,382 pairs and 623,804 singletons were assembled with SPADes version 3.8.2 (10), and contigs that matched either the phiX174 ge-

nome or had less than 100× coverage were excluded. This yielded 27 contigs, with an N_{50} of 250,005 bases, a total length of 3,088,030 bases, an average coverage of 145×, and a GC content of 73%. Gene prediction and annotation were carried out using the RAST pipeline (11) as provided by the PATRIC server (12). A total of 2,905 coding DNA sequences, 45 tRNA genes, and three rRNA genes were identified.

Accession number(s). The nucleotide sequence and annotation data for the *Tessaracoccus lapidicaptus* IPBSL-7^T draft genome have been deposited at DDBJ/ENA/GenBank under the accession number [MBQD00000000](https://www.ncbi.nlm.nih.gov/nuclseq/MBQD00000000/). The version described in this paper is the first version, MBQD01000000.

ACKNOWLEDGMENTS

We thank Victor Parro, Ricardo Amils, and Kenneth N. Timmis, as well as to the other members of the IPBSL project team. We also thank Javier Tamames for his useful advice on genome assembly and annotation.

FUNDING INFORMATION

This work, including the efforts of Alejandro Arce-Rodríguez, was funded by EC | European Research Council (ERC) (ERC250350). This work, including the efforts of Fernando Puente-Sánchez, was funded by Ministerio de Economía y Competitividad (Ministry of Economy and Competitiveness) (CTM2013-48292-C3).

REFERENCES

- Maszenan AM, Seviour RJ, Patel BKC, Schumann P, Rees GN. 1999. *Tessaracoccus bendigoensis* gen. nov., sp. nov., a Gram-positive coccus occurring in regular packages or tetrads, isolated from activated sludge biomass. *Int J Syst Bacteriol* 49:459–468. [http://dx.doi.org/10.1099/00207713-49-2-459](https://doi.org/10.1099/00207713-49-2-459).
- Finster KW, Cockell CS, Voytek MA, Gronstal AL, Kjeldsen KU. 2009. Description of *Tessaracoccus profundus* sp. nov., a deep-subsurface actinobacterium isolated from a Chesapeake impact crater drill core (940 m depth). *Antonie van Leeuwenhoek* 96:515–526. [http://dx.doi.org/10.1007/s10482-009-9367-y](https://doi.org/10.1007/s10482-009-9367-y).
- Puente-Sánchez F, Sánchez-Román M, Amils R, Parro V. 2014. *Tessaracoccus lapidicaptus* sp. nov., an actinobacterium isolated from the deep

- subsurface of the Iberian pyrite belt. *Int J Syst Evol Microbiol* 64: 3546–3552. <http://dx.doi.org/10.1099/ijso.0.060038-0>.
4. Amils R, Fernández-Remolar D, Parro V, Rodríguez-Manfredi JA, Timmis K, Oggerin M, Sánchez-Román M, López JF, Fernández JP, Puente F, Gómez-Ortiz D, Briones C, Gómez F, Omeregíe EO, García M, Rodríguez N, Sanz JL. 2013. Iberian pyrite belt subsurface life (IPBSL), a drilling project of biohydrometallurgical interest. *Adv Mater Res* 825:15–18. <http://dx.doi.org/10.4028/www.scientific.net/AMR.825.15>.
 5. Amils R, Fernández-Remolar D, Parro V, Rodríguez-Manfredi JA, Oggerin M, Sánchez-Román M, López FJ, Fernández-Rodríguez JP, Puente-Sánchez F, Briones C, Prieto-Ballesteros O, Tornos F, Gómez F, García-Villadangos M, Rodríguez N, Omeregíe E, Timmis K, Arce A, Sanz JL, Gómez-Ortiz D. 2014. Río Tinto: a geochemical and mineralogical terrestrial analogue of Mars. *Life* 4:511–534. <http://dx.doi.org/10.3390/life4030511>.
 6. Sánchez-Román M, Puente-Sánchez F, Parro V, Amils R. 2015. Nucleation of Fe-rich phosphates and carbonates on microbial cells and exopolymeric substances. *Front Microbiol* 6:1024. <http://dx.doi.org/10.3389/fmicb.2015.01024>.
 7. Hoehler TM, Jørgensen BB. 2013. Microbial life under extreme energy limitation. *Nat Rev Microbiol* 11:83–94. <http://dx.doi.org/10.1038/nrmicro2939>.
 8. Martin M. 2011. Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnetJ* 17:10–12. <http://dx.doi.org/10.14806/ej.17.1.200>.
 9. Schmieder R, Edwards R. 2011. Quality control and preprocessing of metagenomic datasets. *Bioinformatics* 27:863–864. <http://dx.doi.org/10.1093/bioinformatics/btr026>.
 10. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Pribelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <http://dx.doi.org/10.1089/cmb.2012.0021>.
 11. Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K, Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O, Vonstein V, Wilke A, Zagnitko O. 2008. The RAST server: Rapid Annotations using Subsystems Technology. *BMC Genomics* 9:75. <http://dx.doi.org/10.1186/1471-2164-9-75>.
 12. Wattam AR, Abraham D, Dalay O, Disz TL, Driscoll T, Gabbard JL, Gillespie JJ, Gough R, Hix D, Kenyon R, Machi D, Mao C, Nordberg EK, Olson R, Overbeek R, Pusch GD, Shukla M, Schulman J, Stevens RL, Sullivan DE, Vonstein V, Warren A, Will R, Wilson MJC, Yoo HS, Zhang C, Zhang Y, Sobral BW. 2014. PATRIC, the bacterial bioinformatics database and analysis resource. *Nucleic Acids Res* 42:D581–D591. <http://dx.doi.org/10.1093/nar/gkt1099>.