

Supplementary Material

Host RNA biomarkers for tuberculosis: evaluation of *DOCK9*, *EPHA4*, and *NPC2* expression modulations in blood.

Leonardo Silva de Araujo, Lea A. I. Vaas, Marcelo Ribeiro-Alves, Fernanda Carvalho Queiroz Mello, Alexandre Silva de Almeida, Adriana da Silva Resende Moreira, Afrânio Lineu Kritski, José Roberto Lapa e Silva, Milton Ozório Moraes, Frank Pessler, and Maria Helena Féres Saad.

Corresponding author: Dr. Maria Helena Féres Saad: saad@ioc.fiocruz.br

Supplementary Table S1 - Forward and reverse primers used for the RT-qPCR amplifications.

Sense	Gene	Sequence
Forward	<i>DOCK9</i>	5'-AGAGCCACACAGGAAGAAGTC-3'
	<i>EPHA4</i>	5'-ATGCAGTGGGCGTCTCAAAG-3'
	<i>NPC2</i>	5'-GTCCCAGTTCCTTTCCCAT-3'
	<i>RPL13A</i>	5'-GACAAGAAAAAGCGGATGGT-3'
Reverse	<i>DOCK9</i>	5'-TGGCTCTCCAATCCTTCCTCA-3'
	<i>EPHA4</i>	5'-TGGTTTACATTTAGTGACCACGCC-3'
	<i>NPC2</i>	5'-ACTCCACCACCAGTTTTATAGAGG-3'
	<i>RPL13A</i>	5'-GTA CTCCAGCCAACCTCGT-3'