

Supplementary Material

Host RNA biomarkers for tuberculosis: evaluation of *DOCK9*, *EPHA4*, and *NPC2* expression modulations in blood.

Leonardo Silva de Araujo, Lea A. I. Vaas, Marcelo Ribeiro-Alves, Fernanda Carvalho Queiroz Mello, Alexandre Silva de Almeida, Adriana da Silva Resende Moreira, Afrânio Lineu Kritski, José Roberto Lapa e Silva, Milton Ozório Moraes, Frank Pessler, and Maria Helena Féres Saad.

Corresponding author: Dr. Maria Helena Féres Saad: saad@ioc.fiocruz.br

Supplementary Table S2 - Mean and median normalized expression values of *DOCK9*, *EPHA4*, and *NPC2* mRNAs in the whole blood from the Brazilian cohort.

	n	DOCK9		EPHA4		NPC2	
		Mean	Median	Mean	Median	Mean	Median
G.I	12	0.3527 3	0.0203 4	0.1095 8	0.0819 6	0.0799 6	0.0706 8
G.II	24	0.1768 8	0.0388 0.0337	1.1611 0.4377	0.1298 5	0.8410 9	0.1455 1
G.III	48	0.0955 0.2641	5 0.0444	9 0.9835	9 0.2167	4 1.1880	7 0.3823
G.IV	29	3 2.0720	6 0.0728	2 2.3187	2 0.1718	7 1.9340	6 0.2072
G.V	12	9 0.0299	8 0.0172	4 0.1166	5 0.1041	3 0.1640	2 0.1644
G.VI	4	5	5	5	6	8	3