

Table S3. Geographical dispersal of HBV infections and genotypes

Continent & Region	A	B	C	D	E	F	G	H	I	Total HBV
	(% of global infections with respective genotype)									
AFRICA	73.6	0.05	0.10	22.0	99.4	0.00	0.00	0.00	0.00	35.3
<i>Eastern Africa</i>	44.6	0.05	0.10	10.6	6.79	0.00	0.00	0.00	0.00	11.3
<i>Middle Africa</i>	12.8	0.00	0.00	0.19	13.7	0.00	0.00	0.00	0.00	4.62
<i>Northern Africa</i>	1.35	0.00	0.00	10.1	2.41	0.00	0.00	0.00	0.00	3.02
<i>Southern Africa</i>	7.80	0.00	0.00	0.41	0.45	0.00	0.00	0.00	0.00	1.49
<i>Western Africa</i>	7.01	0.00	0.00	0.76	76.1	0.00	0.00	0.00	0.00	14.9
ASIA	17.2	98.8	98.6	61.9	0.19	0.23	0.00	0.00	100	57.9
<i>Central Asia</i>	0.88	0.00	0.05	6.91	0.00	0.00	0.00	0.00	0.00	1.69
<i>Eastern Asia</i>	0.41	61.0	76.6	5.00	0.00	0.13	0.00	0.00	94.0	30.5
<i>South-Eastern Asia</i>	4.89	5.36	36.1	17.0	0.13	0.00	0.00	0.00	0.00	6.05
<i>Southern Asia</i>	9.46	1.73	4.99	37.2	0.05	0.00	0.00	0.00	0.00	12.1
<i>Western Asia</i>	1.13	0.00	0.03	12.7	0.14	0.10	0.00	0.00	0.00	3.10
EUROPE	3.90	0.17	0.27	13.5	0.14	1.88	9.44	3.25	0.00	4.00
<i>Eastern Europe</i>	1.94	0.01	0.09	9.76	0.00	0.12	0.00	1.93	0.00	2.70
<i>Northern Europe</i>	0.09	0.04	0.02	0.14	0.01	0.01	0.55	0.00	0.00	0.06
<i>Southern Europe</i>	1.08	0.03	0.07	2.74	0.06	1.43	4.23	1.33	0.00	0.85
<i>Western Europe</i>	0.79	0.09	0.09	0.86	0.07	0.32	4.66	0.00	0.00	0.38
LATIN AMERICA/CARIBBEAN	4.51	0.05	0.11	1.96	0.19	97.7	73.5	95.2	0.00	2.20
<i>Caribbean</i>	2.59	0.00	0.00	0.65	0.19	0.00	0.00	0.70	0.00	0.61
<i>Central America</i>	0.09	0.00	0.01	0.03	0.00	4.33	23.8	94.5	0.00	0.13
<i>South America</i>	1.83	0.04	0.10	1.29	0.00	93.4	49.7	0.00	0.00	1.45
NORTHERN AMERICA	0.77	0.83	0.47	0.25	0.04	0.16	17.1	1.58	0.00	0.44
OCEANIA *	0.02	0.14	0.42	0.30	0.00	0.00	0.00	0.00	0.00	0.20

Numbers are rounded to 3 significant digits and represent % of global infections with respective HBV genotype in the indicated world region. * Oceania includes pooled data of Australia/New Zealand, Melanesia, Micronesia, Polynesia