



**HELMHOLTZ  
ZENTRUM FÜR  
INFEKTIONSFORSCHUNG**

**This is an by copyright after embargo allowed publisher's PDF of an article  
published in**

**Elhariry, H.M., Meens, J., Stehr, M., Auling, G.  
S434F in Nrde generates the thermosensitive phenotype  
of Corynebacterium ammoniagenes CH31 and enhances  
thermolability by increasing the surface  
hydrophobicity of the Nrde(Ts) protein  
(2005) Applied and Environmental Microbiology, 71 (9),  
pp. 5582-5586.**

# S434F in NrdE Generates the Thermosensitive Phenotype of *Corynebacterium ammoniagenes* CH31 and Enhances Thermolability by Increasing the Surface Hydrophobicity of the NrdE(Ts) Protein

Hesham M. Elhariry,<sup>1†</sup> Jochen Meens,<sup>1‡</sup> Matthias Stehr,<sup>2</sup> and Georg Auling<sup>1\*</sup>

Institut für Mikrobiologie, Universität Hannover, Schneiderberg 50, D-30167 Hannover, Germany,<sup>1</sup> and GBF—Gesellschaft für Biotechnologische Forschung mbH, Mascheroder Weg 1, D-38124 Braunschweig, Germany<sup>2</sup>

Received 25 November 2004/Accepted 31 March 2005

**The thermosensitive phenotype of strain CH31, a derivative of *Corynebacterium ammoniagenes* ATCC 6872, was allocated by cloning, sequencing, and genetic complementation to a single C→T exchange in the *nrdE* (nucleotide reduction) gene at nucleotide 1301. Protein modeling indicates the impaired surface hydrophobicity of NrdE(Ts) due to the S434F transition.**

A unique enzyme, ribonucleotide reductase (EC 1.17.4.1), delivers the building blocks for DNA replication in all living organisms in a potentially rate-limiting step (9, 29, 34, 38). In the gram-positive nucleotide producer (2, 4, 6) *Corynebacterium* (formerly *Brevibacterium*) *ammoniagenes*, this essential enzyme is a manganese protein (Mn-ribonucleotide reductase) encoded by the *nrdEF* (nucleotide reduction) genes, and the enzymatically active holoenzyme has an  $\alpha\beta_2$  subunit structure (5, 10, 24, 25, 44). By random chemical mutagenesis of *Corynebacterium ammoniagenes* ATCC 6872 using *N*-methyl-*N'*-nitro-*N*-nitrosoguanosine (MNNG), temperature-sensitive (TS) clones which failed to reduce ribonucleotides to 2'-deoxyribonucleotides were obtained (18). Through in vitro complementation, their defect was located biochemically on the CA1 protein, encoded by the *nrdE* gene (18, 24), and therefore renamed R2E (NrdE). Recently, one of these TS mutants, strain CH31, was exploited for nucleotide overproduction by temperature shift in the presence of excess manganese (1).

Here we characterize the *nrdE* gene of *C. ammoniagenes* CH31 with respect to the putative point mutation expected (33) from the foregoing mutagenization with the alkylating compound MNNG.

First, the *nrdE* genes of CH31 and its parental wild type, *C. ammoniagenes* ATCC 6872, were cloned from cetyltrimethylammonium bromide-extracted (19) chromosomal DNA into *Xma*I-digested pUC18 and, after ligation, transformed into *Escherichia coli* XL1-Blue. Strains of *C. ammoniagenes* were routinely cultivated in seed medium (18) at 27°C and *E. coli*

XL1-Blue at 37°C in LB (25) supplemented with ampicillin (100 µg/ml), chloramphenicol (30 µg/ml), D-glucose (0.5%, wt/vol), or IPTG (isopropyl-β-D-thiogalactopyranoside; 1 mM), as required. A 0.5-kb fragment of the *nrdE* gene from *C. ammoniagenes* ATCC 6872 (Fig. 1) was amplified and digoxigenin labeled (16) by PCR using primers 5'-GGCCAGAGAACTCCACGGC-3' (forward) and 5'-TTGTCCATGTGTGGAGCTG-GG-3' (backward) for specific probing (30) of the two chromosomal 5.2-kb *Xma*I fragments and the recombinant plasmids pUCEF6872 and pUCEFCH31 (Table 1). A 2.3-kb region of the *nrd* operon of *C. ammoniagenes* was sequenced (Fig. 1) by a primer walking approach and the chain termination method (31) using a BigDye Terminator cycle sequencing kit on an ABI Prism 310 genetic analyzer (PE Applied Biosystems Inc.). The complete *nrdE* gene, including the intergenic region upstream and 100 bp downstream, was analyzed using the DNASTAR software package (DNASTAR Inc., Madison, WI) and Clone Manager 5.0 (Scientific & Educational Software) and CLUSTAL W for multiple sequence alignments (39). Sequence comparison between the wild type (accession no. AY769914) and mutant (accession no. AY788949) revealed only a single base pair substitution, at position 1301 (C→T), within the *nrdE* coding region, causing a serine-to-phenylalanine exchange at position 434 in NrdE (S434F). Only when we compared the sequence to one deposited *nrdE* gene sequence (accession no. Y09572 [8]) did three additional base

\* Corresponding author. Mailing address: Institut für Mikrobiologie der Universität, Schneiderberg 50, D-30167 Hannover, Germany. Phone: 49/511-762 5241. Fax: 49/511-762 5287. E-mail: auling@ifmb.uni-hannover.de.

† Present address: Department of Food Science, Faculty of Agriculture, Ain Shams University, Cairo, Egypt.

‡ Present address: Zentrum für Infektionsmedizin-Institut für Mikrobiologie, Stiftung Tierärztliche Hochschule Hannover, Bischofsholer Damm 15, D-30173 Hannover, Germany.



FIG. 1. Restriction map of the *nrd* locus of *C. ammoniagenes* strains ATCC 6872 (24, 40) and CH31 (7). The arrow indicates the point mutation within the *nrdE* gene. The position of the *nrdE* hybridization probe is shown as a black bar.

TABLE 1. Microorganisms and plasmids

Microorganism	Related genotype, phenotype, or feature	Source/reference
<b>Strains</b>		
<i>Corynebacterium ammoniagenes</i> Strain CH31	ATCC 6872, wild type, <i>nrdE</i> <sup>+</sup>	Institute's culture collection 18
<i>Escherichia coli</i> XL1-Blue	Thermosensitive mutant of ATCC 6872, <i>nrdE</i> (Ts) <i>endA1 gyrA96 hsdR17</i> ( <i>r<sub>K</sub></i> <sup>-</sup> <i>m<sub>K</sub></i> <sup>+</sup> ) <i>recA1 relA1 supE44 thi-1 F'</i> ( <i>proAB lacI<sup>q</sup>ZΔM15 Tn10</i> )	Stratagene
<b>Plasmids</b>		
pUC18	Amp <sup>r</sup> , cloning vector	45
pUCEF6872	pUC18 with a 5.2-kb XmaI fragment from <i>C. ammoniagenes</i> ATCC 6872	This work
pUCEFCH31	pUC18 with a 5.2-kb XmaI fragment from <i>C. ammoniagenes</i> CH31	This work
pXMJ19	Cm <sup>r</sup> , <i>C. glutamicum</i> / <i>E. coli</i> shuttle vector	13
pXE6872	pXMJ19 containing the <i>nrdE</i> <sup>+</sup> gene from <i>C. ammoniagenes</i> ATCC 6878 (accession no. AY769914) cloned by PCR	This work
pXECH31	pXMJ19 containing the <i>nrdE</i> (Ts) gene from <i>C. ammoniagenes</i> CH31 (accession no. AY788949) cloned by PCR	This work

pair substitutions appear within a stretch of only 43 bp at nucleotide (nt) 2021 (G→A), nt 2032 (G→A), and nt 2063 (T→A). However, we consider the clustered amino acid differences at positions 674, 678, and 688, showing up only upon comparison to the NrdE sequence published in GenBank (CAA70765, gi:3077613), as irrelevant due to their absence in four other corynebacterial and five mycobacterial NrdE sequences (<http://www.ncbi.nih.gov/entrez>).

For recombinant expression, both the *nrdE*<sup>+</sup> and *nrdE*(Ts) genes were subcloned without any flanking regions but including their ribosome binding sites from the two sequenced plasmids, pUCEF6872 and pUCEFCH31, using the primers 5'-G GGGTCTAGATTGAAAGGCCGAGTGCTTCAAATGAC-3' (forward) and 5'-AAAGGAGCTCTTAGAGCATGCGGAGACGCAACC-3' (backward). The additional XbaI site and SacI site are underlined. The start codon was changed from GTG to ATG by the forward primer. The XbaI/SacI-digested amplicons were purified prior to ligation into the XbaI/SacI-digested *E. coli*/*C. glutamicum* shuttle vector pXMJ19 (13). The fact that the resulting expression vectors, pXE6872 (*nrdE*) and pXECH31 [*nrdE*(Ts)], were first introduced into *E. coli* XL1-Blue and NrdE synthesis from the plasmid constructs was proven by immunostaining with anti-NrdE antibodies (not shown). The entire nucleotide sequence of the *nrdE*<sup>+</sup> and *nrdE*(Ts) genes was confirmed by double-

strand sequencing. Then, the plasmids pXMJ19, pXE6872, and pXECH31 were introduced into the TS mutant, *C. ammoniagenes* CH31, by electroporation (40) with a BTX Electro Cell Manipulator ECM600. Only the transformant CH31/pXE6872 recovered and displayed wild-type morphology at 37°C (not shown). Expression of the introduced *nrdE*<sup>+</sup> gene and formation of fully functional holoenzyme in liquid culture (not shown) required an induction period of 4 hours prior to the temperature shift. On NBH agar (8 g nutrient broth, 2 g yeast extract, and 5 g NaCl per liter) with 1 M IPTG added, the growth defect of the TS mutant CH31 was complemented by plasmid pXE6872, whereas pXECH31 and the shuttle vector




FIG. 2. Complementation of the growth defect of the TS mutant CH31 carrying the empty vector pXMJ19 (A), plasmid pXECH31 (B), or plasmid pXE6872 (C) after 3 days on NBH agar supplemented with 1 mM IPTG.




FIG. 3. Expression of *nrdE*<sup>+</sup> and *nrdE*(Ts) in *C. ammoniagenes* CH31 shown by Coomassie blue-stained sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) (15) of whole-cell proteins (a) or a Western blot (41) probed with NrdE-specific antibodies (b). Molecular weight markers (Amersham Pharmacia Biotech) were run in lane M. Molecular weights (in thousands) are noted at the left. Extracts from transformants of CH31 carrying plasmids pXMJ19 are seen in lanes 1 and 2, pXE6872 in lanes 3 and 4, and pXECH31 in lanes 5 and 6. For a 4-h induction period, cells were grown in NBH broth at 27°C in the absence (lanes 1, 3, and 5) or presence (lanes 2, 4, and 6) of 1 mM IPTG. The arrows indicate the bands of NrdE protein which were synthesized only upon induction. Cellular proteins were obtained from 2-ml cultures of washed *C. ammoniagenes* cells after incubation in 100 μl lysis buffer (10 mM Tris-HCl, pH 6.8, 25 mM MgCl<sub>2</sub>, 200 mM NaCl), containing 5 mg/ml lysozyme, for 60 min at 37°C and subsequent addition of 10 μl SDS (10%) and 100 μl loading buffer (15) prior to being heated at 95°C for 5 min.



were not effective (Fig. 2). The failure of pXECH31 was not due to inefficient expression of the introduced *nrde*(Ts) gene since CH31/pXE6872 and CH31/pXECH31 produced the same amount of NrdE (Fig. 3) upon IPTG induction.

The finding of the nonconservative S434F exchange suggested a location for this point mutation in the tertiary structure and generated a model (720 amino acids, residues 17 to 705) of *C. ammoniagenes* NrdE (Fig. 4a) using the structure (42) of *Salmonella enterica* serovar Typhimurium NrdE (PDB code 1PEO, 68% sequence identity, r.m.s of 0.176 Å with 679 equivalent residues). Serine 434 is distant from the active (30-Å) and effector binding (60-Å) sites and located at a surface-exposed loop, connecting helix  $\alpha$ 14 with the following helix, as a very conserved residue in a structural motif, containing polar and charged residues, S434, R161, (D/E)436, and (T/S)440 common to NrdE proteins (Fig. 5). Conservation of surface hydroxyl groups in NrdE proteins may reflect a requirement for hydrogen bonding with the solvent. A hypothetical water molecule, crucial for thermostability, also present at conserved positions of other proteins (35, 36), stabilizes the surface-exposed location of Ser434 by an extensive network of hydrogen bonds (Fig. 4B). This network collapses by displacement of the central water molecule resulting from the S434F mutation, and the surface loop is destabilized (Fig. 4C). Strikingly similar properties (distance from the active site, surface exposition, lack of intramolecular hydrogen bonding) are shown by the T157I mutant, the most thermolabile variant in a series of T4 lysozyme mutants, pointing to increased thermostability resulting from increased surface hydrophobicity (3, 11, 20), while relief of reverse hydrophobic effects generates increased thermostability (28, 37, 43).

Further analysis of the S434 surface loop using the expression system developed here is suggested since neither the degradation of NrdE proteins nor leakiness of the strong *tac* promoter (this work and reference 27) occurred in *C. ammoniagenes*. Beyond this, our TS mutant CH31 may serve as a safe host strain for functional studies of foreign *nrde* genes (e.g., from mycobacteria) restricted in the original pathogenic background, since it did not revert in the decade following its isolation.

We are grateful to A. Burkovski for providing the *C. glutamicum*/*E. coli* shuttle vector plasmid pXMJ19. G. Auling thanks H. Diekmann for stimulating discussions and continued interest in perturbations of the corynebacterial cell cycle.

H. M. Elhariry thanks the Ministry of Higher Education and Scientific Research, Arabic Republic of Egypt, for a long-term fellowship in support of his Ph.D. thesis.

#### REFERENCES

1. Abbouni, B., H. M. Elhariry, and G. Auling. 2004. Overproduction of NAD<sup>+</sup> and 5'-inosine monophosphate in the presence of 10  $\mu$ M Mn<sup>2+</sup> by a mutant of *Corynebacterium ammoniagenes* with thermosensitive nucleotide reduction (*nrde*<sup>ts</sup>) after temperature shift. *Arch. Microbiol.* **182**:119–125.
2. Abbouni, B., H. M. Elhariry, and G. Auling. 2003. Arrest of cell cycle by inhibition of ribonucleotide reductase induces accumulation of NAD<sup>+</sup> by Mn<sup>2+</sup> supplemented growth of *Corynebacterium ammoniagenes*. *Biotechnol. Lett.* **25**:143–147.
3. Alber, T., D. P. Sun, K. Wilson, J. A. Wozniak, S. P. Cook, and B. W. Matthews. 1987. Contributions of hydrogen bonds of Thr 157 to the thermodynamic stability of phage T4 lysozyme. *Nature* **330**:41–46.
4. Auling, G., and H. Follmann. 1994. Manganese-dependent ribonucleotide reduction and overproduction of nucleotides in coryneform bacteria. *Metal Ions Biol. Syst.* **30**:131–164.
5. Auling, G., M. Thaler, and H. Diekmann. 1980. Parameters of unbalanced growth and reversible inhibition of deoxyribonucleic acid synthesis in *Brevibacterium ammoniagenes* ATCC 6872 induced by depletion of Mn<sup>2+</sup>. Inhibitor studies on the reversibility of deoxyribonucleic acid synthesis. *Arch. Microbiol.* **127**:105–114.
6. Elhariry, H., H. Kawasaki, and G. Auling. 2004. Recent advances in microbial production of flavour enhancers for the food industry. *Recent Res. Dev. Microbiol.* **8**:15–39.
7. Elhariry, H. M. 2004. Characterization of a thermosensitive ribonucleotide reductase mutant derived from *Corynebacterium ammoniagenes* ATCC 6872 and its use in the production of ribonucleotides. Ph.D. thesis. University of Hannover, Hannover, Germany.
8. Fieschi, F., E. Torrents, L. Touloukhonova, A. Jordan, U. Hellmann, J. Barbe, I. Gibert, M. Karlsson, and B.-M. Sjöberg. 1998. The manganese-containing ribonucleotide reductase of *Corynebacterium ammoniagenes* is a class Ib enzyme. *J. Biol. Chem.* **273**:4329–4337.
9. Follmann, H. 2004. Deoxyribonucleotides: the unusual chemistry and biochemistry of DNA precursors. *Chem. Soc. Rev.* **33**:225–233.
10. Griepenburg, U., R. Kappl, J. Hüttermann, and G. Auling. 1998. A divalent metal site in the small subunit of the manganese-dependent ribonucleotide reductase of *Corynebacterium ammoniagenes*. *Biochemistry* **37**:7992–7996.
11. Grütter, M. G., T. M. Gray, L. H. Weaver, T. A. Wilson, and B. W. Matthews. 1987. Structural studies of mutants of the lysozyme of bacteriophage T4. The temperature-sensitive mutant protein Thr157→Ile. *J. Mol. Biol.* **197**:315–329.
12. Guex, N., and M. C. Peitsch. 1997. Protein modelling via the internet. *BIO World* **5**:9–13.
13. Jakoby, M., C. E. Nguoto-Nkii, and A. Burkovski. 1999. Construction and application of new *Corynebacterium glutamicum* vectors. *Biotechnol. Tech.* **13**:437–441.
14. Kraulis, P. J. 1991. MOLSCRIPT: a program to produce both detailed and schematic plots of protein structures. *J. Appl. Crystallogr.* **24**:946–950.
15. Laemmli, U. K. 1970. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* **227**:680–685.
16. Lanzillo, J. J. 1990. Preparation of digoxigenin-labeled probes by the polymerase chain reaction. *Biotechnology* **8**:620–622.
17. Laskowski, R. A., M. W. MacArthur, D. S. Moss, and J. M. Thornton. 1993. Procheck: a program to check stereochemical quality of protein structures. *J. Appl. Crystallogr.* **26**:283–290.
18. Luo, C. H., J. Hansen, and G. Auling. 1997. Temperature-sensitive mutants of *Corynebacterium ammoniagenes* ATCC 6872 with a defective large subunit of the manganese-containing ribonucleotide reductase. *Arch. Microbiol.* **167**:317–324.
19. Martin, J. F., and J. A. Gil. 1999. *Corynebacteria*, p. 379–391. In A. L. Demain and J. E. Davies (ed.), *Manual of industrial microbiology and biotechnology*, 2nd ed. ASM Press, Washington, D.C.
20. Matsumura, M., Y. Katakura, T. Imanaka, and S. Aiba. 1984. Enzymatic and nucleotide sequence studies of a kanamycin-inactivating enzyme encoded by a plasmid from thermophilic bacilli in comparison with that encoded by plasmid pUB110. *J. Bacteriol.* **160**:413–420.
21. Merritt, E. A., and D. J. Bacon. 1997. Raster3D: photorealistic molecular graphics. *Methods Enzymol.* **277**:505–524.
22. Miller, J. H. 1972. *Experiments in molecular genetics*. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.
23. Nicholas, K. B., H. B. Nicholas, and D. W. Deerfield. 1997. GeneDoc: analysis and visualization of genetic variation. *EMBNETNEWS* **4**, 2.
24. Oehlmann, W. 1998. Klonierung der Gene der Ribonucleotid-Reduktasen von *Corynebacterium ammoniagenes* und *Corynebacterium glutamicum*. Ph.D. thesis, University of Hannover, Hannover, Germany.
25. Oehlmann, W., U. Griepenburg, and G. Auling. 1998. Cloning and sequencing of the *nrde* gene of *Corynebacterium ammoniagenes* ATCC 6872 encoding the functional metallo-cofactor of the manganese-ribonucleotide reductase (Mn-RNR). *Biotechnol. Lett.* **20**:483–488.
26. Orengo, C. A., A. D. Michie, S. Jones, D. T. Jones, M. B. Swindells, and J. M. Thornton. 1997. CATH—a hierarchic classification of protein domain structures. *Structure* **5**:1093–1108.
27. Paik, J. E., and B. R. Lee. 2003. Isolation of transcription initiation signals from *Corynebacterium ammoniagenes* and comparison of their gene expression levels in *C. ammoniagenes* and *Escherichia coli*. *Biotechnol. Lett.* **25**:1311–1316.
28. Pakula, A. A., and R. T. Sauer. 1990. Reverse hydrophobic effects relieved by amino-acid substitutions at a protein surface. *Nature* **344**:363–364.
29. Reichard, P. 1993. From RNA to DNA. Why so many ribonucleotide reductases? *Science* **260**:1773–1777.
30. Sambrook, J., E. F. Fritsch, and T. Maniatis. 1989. *Molecular cloning: a laboratory manual*, 2nd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.
31. Sanger, F., S. Nicklen, and A. R. Coulson. 1977. DNA sequencing with chain-terminating inhibitors. *Proc. Natl. Acad. Sci. USA* **74**:5463–5467.
32. Schwede, T., J. Kopp, N. Guex, and M. C. Peitsch. 2003. SWISS-MODEL: an automated protein homology-modeling server. *Nucleic Acids Res.* **31**:3381–3385.
33. Seiffert, W. 2003. *Lehrbuch der Genetik*, 2nd ed. Spektrum Akademischer Verlag, Heidelberg, Germany.

34. **Sjöberg, B.-M.** 1997. Ribonucleotide reductases—a group of enzymes with different metallosites. *Struct. Bonding* **88**:139–173.
35. **Stehr, M., G. Schneider, F. Åslund, A. Holmgren, and Y. Lindqvist.** 2001. Structural basis for the thioredoxin-like activity profile of the glutaredoxin-like NrdH-redoxin from *Escherichia coli*. *J. Biol. Chem.* **276**:35836–35841.
36. **Stehr, M., and Y. Lindqvist.** 2004. NrdH-redoxin of *Corynebacterium ammoniagenes* forms a domain-swapped dimer. *Proteins* **55**:613–619.
37. **Strub, C., C. Alies, A. Lougarre, C. Ladurantie, J. Czaplicki, and D. Fournier.** 2004. Mutation of exposed hydrophobic amino acids to arginine to increase protein stability. *BMC Biochem.* **5**:9.
38. **Stubbe, J., and W. van der Donk.** 1995. Ribonucleotide reductase: radical enzymes with suicidal tendencies. *Chem. Biol.* **2**:793–801.
39. **Thompson, J. D., T. J. Gibson, F. Plewniak, F. Jeanmougin, and D. G. Higgins.** 1997. The CLUSTAL X windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Res.* **25**:4876–4882.
40. **Torrents, E., I. Roca, and I. Gibert.** 2003. *Corynebacterium ammoniagenes* class Ib ribonucleotide reductase: transcriptional regulation of an atypical genomic organization in the *nrd* cluster. *Microbiology* **149**:1011–1020.
41. **Towbin, H. G., T. Staehelin, and J. Gordon.** 1979. Electrophoretic transfer of proteins from polyacrylamide gels to nitrocellulose sheets: procedure and some applications. *Proc. Natl. Acad. Sci. USA* **76**:4350–4354.
42. **Uppsten, M., M. Färnegårdh, A. Jordan, R. Eliasson, H. Eklund, and U. Uhlin.** 2003. Structure of the large subunit of class Ib ribonucleotide reductase from *Salmonella typhimurium* and its complexes with allosteric effectors. *J. Mol. Biol.* **330**:87–97.
43. **Vogt, G., S. Woell, and P. Argos.** 1997. Protein thermal stability, hydrogen bonds, and ion pairs. *J. Mol. Biol.* **269**:631–643.
44. **Willing, A., H. Follmann, and G. Auling.** 1988. Ribonucleotide reductase of *Brevibacterium ammoniagenes* is a manganese enzyme. *Eur. J. Biochem.* **170**:603–611.
45. **Yanisch-Perron, C., J. Vieira, and J. Messing.** 1985. Improved M13 vectors and host strains: nucleotide sequences of M13mp18 and pUC vectors. *Gene* **33**:103–119.